

Review article

Human dermatosis caused by vesicating beetle products (Insecta), cantharidin and paederin: An overview

Karem S. Ghoneim

Faculty of Science, Al-Azhar University, Cairo, Egypt

Correspondence: Kar_ghoneim@yahoo.com

Abstract

Three major families in the order Coleoptera have been found to produce vesicant toxins, Meloidae (true blister beetles) and Oedemeridae (false blister beetles) produce cantharidin and Staphylinidae (rove beetles) produce paederin. Some of the most important distribution, biological and ecological aspects of these beetle families had been studied because of their products cause human dermatitis. The detection, biosynthesis and chemistry of both cantharidin and paederin were discussed. The prevalence of Paederus dermatitis (skin blistering inflammation) in the world was reviewed. Also, the diagnostic characterization, clinical symptoms of dermatitis and conjunctivitis, the mode of action and pharmacological properties of each vesicant product, preventive measures and curative treatments of the human dermatitis had been included in the present review. Copyright © WJMMMS, all right reserved.

Keywords: biosynthesis, conjunctivitis, endosymbiont, Meloidae, Oedemeridae, Staphylinidae, toxicity, symptom.

Introduction

The order Coleoptera constitutes almost 25% of all known life-forms [1]. The available literature contains information on the development, biology, morphology, epidemiology and control of Coleoptera of medical importance [2]. Ill-effects of Coleoptera range from the immediate trauma of a bite to the effects of the vesicating beetles and also the acute corneal erosion that is attributed to the small beetle *Orthoperus* sp. (family Corylophidae). Other effects are, also, known as a result of beetle metabolites or structures, such as "carpet beetle dermatitis" from the irritating hairs of *Anthrenus* larvae (family Dermestidae), and inhalational asthma from beetles, notably the grain weevil *Sitophilus*, the causative agent of certain cases of grain-worker's asthma. The medical significance of the vesicating darkling beetles (family Tenebrionidae) has been reported [3-6]. The human dermatitis was, also, reported for the tiger beetles (family Cicindelidae) but the mechanism by which the beetles provoke skin symptoms is still a matter of debate [7]. Also, tiger beetles had been reported to cause ocular damage and conjunctiva [8]. In addition to these beetle families, contact with other beetles of family Dermestidae can produce a variety of skin disorders such as dermatitis, vesicular, pustular and vasculitic lesions. Also, there may be pruritus, desquamation and papular urticaria [9, 10].

In Coleoptera, only Meloidae, Oedemeridae and Staphylinidae are the beetles releasing vesicant known chemicals, the first two release cantharidin and the latter releases paederin [11]. Cantharidin is among the most widely known insect natural products [12, 13]. It usually attracted a great attention of many investigators and research institutions because of its toxicosis for man and painful or fatal diseases for horses and other livestock. Many reports on different aspects of

veterinary and medical impacts and uses of cantharidin are available in the literature [14-25]. Paederin is a vesicating agent found in the haemolymph of the *Paederus* genus. It differs from cantharidin, in its chemical and biologic properties [26]. The paederin manufacture is largely confined to adult female beetles. Larvae and males only store paederin acquired maternally (i.e., through eggs) or by ingestion [27].

Human dermatitis is widely reported in the world after contact with the coleopterans in three major families, Meloidae, Oedemeridae and Staphylinidae [28-30]. Bullous lesions are produced when the active ingredient cantharidin is released at the time the beetle is crushed or rubbed upon the exposed skin within hours [31, 32]. Cantharidin can cause severe skin blisters, especially when the insects discharge it from their junctions as a defense measure or get crushed on a human body [33, 34]. Cantharidin in both sexes of *Oedemera podagrariae ventralis* (Oedemeridae) was found in an average of 3.89 µg/beetle in males and 21.68 µg/beetle in females, which are amounts sufficient to irritate human skin [35]. Christmas et al. [36] described seventy-four New Zealand Army personnel with a distinctive bullous dermatosis caused by an endemic beetle *Thelyphassa lineata* (Fabricius) (Oedemeridae) as the first reported cases. Paederus dermatitis (PD) is a blistering inflammation caused when the *Paederus* beetles are accidentally crushed on the human skin [37].

The present review primarily deals with the human skin dermatosis caused by cantharidin or paederin and secondarily is interested in the major three families producing these vesicating chemicals, Meloidae, Oedemeridae and Staphylinidae.

1. Major Beetle families responsible for human dermatosis

1.1. Cantharidin-producing beetles

Family Meloidae (true blister beetles) is virtually cosmopolitan but absent only from New Zealand, Antarctica and most Polynesian islands [38-43]. They primarily occur in temperate steppic and arid regions, and in sub-tropical and tropical savannas or other open habitats [44]. Different biological aspects of this family had been reviewed by Ghoneim [45] as well as various agronomic and biodiversity impacts of these beetles had been reviewed by Ghoneim [46]. These beetles are commonly known as "oil beetles" because the adults release yellow oily droplets of haemolymph from their leg joints (and may be from the antennal joints) when disturbed. This exudation contains toxic material "cantharidin" or "cantharid" which acts naturally as an aphrodisiac for adult males and females (for review, see Ghoneim, 47). It is surprising that the larvae manufacture and accumulate cantharidin as they feed and grow in size. The first five larval instars of the ebony margined blister beetle *Epicauta funebris* Horn were found to produce cantharidin and when disturbed they exude it, as a defensive agent, in a milky oral fluid, not in haemolymph which adult beetles reflexively discharge [48].

Oedemeridae (false blister beetles) is a family of worldwide distribution. It consists of about 1500 species in 100 genera, mostly associated with rotting wood as larvae, though adults are quite common on flowers [49]. They derive their common name not only from their superficial resemblance to some species of Meloidae but also because many species possess the blistering agent cantharidin. Although they are commonly known as false blister beetles, some recent authorities have coined the name pollen-feeding beetles [49]. The available literature contains several reported works on the distribution and systematics of the family Oedemeridae in the Old World regions and countries, such as Russia [50], southeastern China [51], Japan and Taiwan [52], Korea [53], Iran [35], southern Africa [54,55], tropical Africa [56], Socotra Archipelago (in the northwest Indian Ocean)[57], Iberian peninsula [58], eastern Mediterranean region [59], Greece [60], Latvia [61], Bulgaria and Turkey [62, 63], Sweden [64], and some other regions [65-68]. With regard to the distribution and systematics of this family in the New World, many published works had been available in the literature such as Campbell [69], Kriska [70], Majka and Langor [71], Webster et al. [72] for Canada; Blackwelder [73], Lawrence and Newton [74] and Patrice et al. [75] for Central America and South America; Paula [76] for New Zealand; and Arnett [77] for some other regions and countries.

Cantharidin detection and quantitative determination in different species of Oedemeridae had been studied [35, 78-81]. Since most adults of Oedemeridae are soft-bodied flower feeders it is thought that they employ the cantharidin to deter potential predators [70]. Although there are few references in the medical literature, blister beetle dermatosis caused by oedemerids may be more common and widespread than currently recognized [26]. From a public health standpoint,

some of the blister beetles swarm during certain seasons and the large numbers of them pose a considerable threat [82] because they exudate droplets of haemolymph which produce blistering on contact with human skin [15].

1.2. Paederin-producing beetles

The family Staphylinidae (rove beetles) currently comprises 3,847 genera and 56,768 species throughout the world [83]. The distribution and systematics of Staphylinidae had been studied in different parts of the Old World such as China [84, 85], Turkey [86, 87], South Africa [88], Madagascar [89, 90] and Australia [91]. On the other hand, some works had been carried out on the distribution and/or systematics of Staphylinidae in several parts of the New World [92-99]. Staphylinid beetles are completely harmless though their habits make them unappealing. They are found in or near decaying organic matter, especially dead animals. They have the interesting (though unpleasant) habit of feeding on other insects such as fly maggots that infest carrion (less often dung or fungus) [100]. Dekeirsschieter et al. [101] reported the monitoring of the presence of adult rove beetles on decaying pig carcasses during early spring in urban, forest and agricultural biotopes of Western Europe.

From the medical point of view, only three different genera can cause PD: *Paederus*, *Paederidus*, and *Megalopaederus* [102]. Although the genus *Paederus* includes several hundred species, it is believed that only 20-30 of them can cause dermatitis [104]. The genus *Paederus* is widely distributed in the world, especially in all temperate and tropical continents, north and south of the equator [105]. There are no species of *Paederus* in Antarctica [106]. The genus *Paederus* includes more than 622 species [103, 107-109] and is represented in the Palaearctic region with approximately 85 species or subspecies [110]. Among the species living under moist conditions, some have attained their life cycle according to seasonal variations [111]. Temperate-zone species may have a single annual breeding season in the warmer months, but in tropical species timing of the breeding season appears to depend upon rainfall seasonality. Widely distributed species with a broad latitudinal range seem to be multivoltine, with more generations at lower latitudes [106]. Species living in temperate regions may have a single breeding season only in the hot months, but in tropical areas the breeding season depends upon rainfall. The faunal diversity of this genus, especially in the Iranian fauna, was reviewed [24]. The adult beetle does not bite or sting, but accidental brushing against it or crushing it over the human skin provokes the release of its coelomic fluid, which contains a strong blistering chemical, paederin [112].

Paederus beetles are usually bicolourous, black or blue and red, sometimes entire body is black or blue or even reddish. Because of the medical and predatory importance of *Paederus*, the life history had been studied. In some studies of *Paederus*, life history had been conducted to know its predatory potentials by using different pests as a prey [106] and some researchers studied its life history by providing artificial diets [113]. Nasir and Akram [114] studied its life history by providing natural conditions (aphids as food) in the laboratory. In the field, eggs are laid on moist substrate singly and are in danger of desiccation. The larval instars are predatory as do adults in moist habitats. A female laid more number of eggs in moist soil rich in organic matter but very few in dry sandy and clay soil [115]. As reported by Laba and Kilin [116], the larval and egg stages of *Paederus fuscipes* Curtis were 9.2 days and 3.8 days, respectively. The adult longevity was 113.8 days for females and 109.2 days for males. The total number of eggs laid was 106 eggs. Incubation period of egg was 4 days. In the laboratory, eggs hatch at about 14 days. Larvae are whitish with black markings [117]. Pupation occurs in earthen cells [34, 118, 119].

The presence of large numbers of the rove beetle *Paederus affieri* Kock in Egypt, due to the less use of insecticides for cotton pests, is the cause of PD [120] and this beetle flies to the residential buildings being attracted by bright lights. However, it cannot fly for a long distance unless being disturbed by a provocative like smoke arising from the burning dry remains of rice in the intermediate area of cultivable land [121].

The rove beetle *P. fuscipes* is commonly known as the 'Beetle Tomcat' because its shape looks like an American fighter aircraft, TOMCAT. It is common and widely distributed insect throughout the world, except America, since it is recorded in Europe, North and central Africa, Asia, New Guinea, Australia, India, Pakistan, etc. In Italy, *P. fuscipes* lives in the cultivated maize fields, hence the common use of the term "the maize field wildfire" to indicate the dermatitis provoked by this beetle [122, 123]. Some bioecological aspects of *P. fuscipes* were studied in the north-west parts of Mazandaran province (Iran) by Abbasipour [124]. It appears in May and lasted until October and they spent 6-8 months in hibernation according to different localities. The daily and seasonal activities were recorded. The seasonal

outbreak of PD, in Permatang Pauh, Malaysia had been described [125]. Huge paddy field areas and high amount of rainfall are the main factors increasing *P. fuscipes* density and PD cases. In Malaysia, also, Bong et al. [126] investigated the life history characteristics of *P. fuscipes* under laboratory conditions using three field strains. The total developmental time of immature stages differed significantly among the three strains. Nasir et al. [127] investigated and related *P. fuscipes* assemblages to cultivated soils and forest parameters of the Punjab, Pakistan during 2008-2009. It was mainly recorded from maize (*Zea mays*) and clover (*Trifolium alexandrinum*) crops due to presence of soft bodied insects. It preferred damp soil rich in organic matter for egg laying under natural conditions. As reported by Caroline [100], the net reproduction rates ranged from 40.09 ± 7.39 to 42.34 ± 8.25 offspring. The rove beetles *Paederus ilvae* (Bernhauer) and *Paederus iliensis* (Coiffait) were identified in the central and western parts of Fars province, Iran. The adults have 5 months activity in average, begin lately in April and then continue until early September; soon afterwards they endure the inclement climate. This over wintering period takes 6-8 month long according to the region [128].

With regard to the ecological and behavioural patterns of *Paederus* beetles, the adults and larvae often live in damp environments [129]. Unlike most staphylinid adults, which avoid daylight, *Paederus* adults are active in the open in daylight. However, flight seems largely restricted to nights of high temperature. Adults of some species are wingless, and in other species adults show individual variation in wing length [106]. The population of *P. fuscipes* was observed under the heaps of leaves to conserve temperature and to keep the habitat wet [130]. In Punjab (Pakistan), *P. fuscipes* was found of its maximum population from grasses near water, from vegetable land and from those crops which had soft bodied insect pests like aphids, mites, leaf hoppers on them [124, 127, 130]. The results of a field study indicated that there was a positive correlation between decaying organic matter (especially dung), soil moisture contents and *P. fuscipes*'s population. This difference was due to biotic factors such as different crops and abiotic factors such as use of pesticides, temperature and soil moisture contents [131, 132]. Seasons also affect the population, e.g. by rain fall [34]. The seasonal effect of moonlight on the vertical distribution of *P. Alfierii* was studied in Qena (Egypt) during two years (1982-83). Moonlight had no significant influence on flight activity in any one season at 1.5 m and at 18.5 m its influence was negligible, even when both years were combined [133].

The *Paederus* adults are active during the day and are attracted by incandescent and fluorescent lights at night, which can bring them into contact with humans [106, 134, 135], especially in tropics and some temperate countries [114]. Black ray was most attractive to the adult beetles and they showed highest daily activities between 21-22 hours at night in Iran [124]. The fluorescent lamp usages at night and accommodations that permit *P. fuscipes* entrance into buildings had contributed to PD incidence in Permatang Pauh, Malaysia [125]. Light had attractive effect on *P. fuscipes*, so maximum activity was observed in the Punjab (Pakistan) during night. Its activity was greatly influenced by light and soil moisture contents [127].

1.3. Beneficial *Paederus* beetles in agriculture and medicine:

It may be important to mention that *Paederus* beetles are beneficial insects because they are carnivorous and eats smaller pests. Large populations of *Paederus* spp. had been recorded from agricultural habitats which make them beneficial due to their feeding on insect pests of major crops, particularly insect pests like the rice moth *Corcyra cephalonica* (Stainton) (Lepidoptera: Galleriidae), the cotton bollworm *Helicoverpa* (=*Heliothis*) *armigera* (Lepidoptera: Noctuidae), the cotton aphid *Aphis gossypii* Glover (Homoptera: Aphididae), the spotted bollworm *Earias vittella* (Fabricius) (Lepidoptera: Noctuidae), the cotton leafworm *Spodoptera litura* (Fabricius) (Lepidoptera: Noctuidae), the rice leaffolder *Marasmia patnalis* Bradley (Lepidoptera: Pyralidae), the soybean aphid *Aphis glycines* Matsumura (Homoptera: Aphididae), fruitfly and many other dipterous and other arthropods [119, 135, 136].

As natural predators, *Paederus fusca* is well known in the rice fields of Southeast Asia, *Paederus crebipunctatis* and *Paederus sabaeus* Erichson occur in East Africa, as well as *P. cruenticollis* is important in Australia. In Indonesia, naturally occurring populations of *P. fuscipes* play an important role in suppressing the soybean aphid. As many as twenty-two different species of prey were predated upon by the predator *P. fuscipes* when offered in the laboratory [137]. In Indonesia, also, *P. fuscipes* is an important predator in soybean fields, including toward *Helimverpa armigera* [138]. The latter authors indicated that population of this natural predator in soybean field were very complex and potential to be used for control the pest populations. In New Delhi (India), the bionomics, predatory potential, field

population levels and seasonal incidence of the predator *P. fuscipes* were studied. This predator had a feeding potential of 0.6 rice leaf folder larvae per individual per day and thus it is an effective predator [139]. Liu and Zheng [140] conducted a study on the predation of *P. fuscipes* on the turnip aphid *Lipaphis erysimi* (Kaltenbach)(Homoptera: Aphididae) and interference with each other. When *P. fuscipes* density grows up, its predation to *L. erysimi* is going to decline. They affect each other. Padmavathi et al. [141] investigated the prevalence, prey preference and predatory potential of *P. fuscipes* in rice. This predator preferred brown planthopper and white backed planthopper nymphs to other prey species tested. In Egypt, Tawfik et al. [142] reported the rove beetle *P. alfierii* as common insects in the fields of maize, cotton and clover. It has a great agricultural benefit as a predator of aphids. The occurrence of *P. alfierii* adult beetles in maize, cotton and clover fields in Giza seemed to follow a certain seasonal succession. Such succession appears to be closely related to the seasonal abundance of their different preys on these crops [143]. Shebl et al. [144] found *P. alfierii* as one of the most common natural enemies associated with pests in alfalfa fields in some parts of Egypt. Field data revealed that *P. alfierii* was found as one of the common insect predators in the fields of Egyptian cotton [145, 146] and of soybean fields [147]. Also, *P. alfierii* was observed among the natural predators of soft scale insects in Egypt [148, 149]. *P. sabaeus* was among the natural predators recommended for controlling the rice yellow mottle virus vectors in some sites of northern Cameron [150]. Thus *Paederus* beetles play an important role as a biological control of insect pests [100]. In addition to its role as a biological control agent, *P. fuscipes* decomposes the organic matter in the soil and hence increases the fertility of soil through recycling of nutrients [127].

Influenced predatory potential of *Paederus* spp. by some of the commonly used insecticides has attracted the attention of some researchers in different parts in the world. Yoshisato [151] evaluated the contact toxicity of several insecticides against *P. fuscipes* under laboratory conditions. It was observed that lindane was most toxic and then DDT. Beta-cyfluthrin was found to be the least toxic to *P. alfierii* and other tested natural enemies of the silverleaf white fly *Bemisia tabaci* (Gennadius) (Homoptera: Aleyrodidae) on cotton in Assiut (Egypt) [152]. A laboratory experiment was conducted by Devi et al. [153] to evaluate the toxicity and residual toxicity of some common insecticides against *P. fuscipes* adults. Their results revealed that neemarin and biorin did not show any toxicity whereas cypermethrin, endosulfan, metasystox and fenvalerate proved highly toxic to this predator. Under laboratory conditions, Meng and Zheng [154] studied the mortalities and effects of bisultap on the predatory function of *P. fuscipes*. The experimental results indicated that the pesticide bisultap has less effect on mortality and predatory function of *P. fuscipes*, but the effects on predatory function of survival individuals are stronger than that on lethality. Using the soaking technique under laboratory conditions, Wang [155] determined the inflict casualties of five kinds of common pesticides on *P. fuscipes*. The results of bioassay showed that: toxicity of 10% chlorpyrifos plus deltamethrin was the strongest, 40% omethoate was the second strongest, 22% imidacloprid plus buprofezin and 50%phoxim plus fenvalerate were comparatively weak, and that of 18% bisultap aqueous solution was the weakest.

From the medical point of view, paederin (the *Paederus* secretion as will be discussed later on) has suppressed cancerous tumors in mice, rats, and plants, has stimulated regeneration of damaged tissues, has healed chronic, necrotic lesions in geriatric patients, and has induced cell fusion in human skin fibroblasts [106]. Paederin is important in cancer and cellular biology [156]. It is worth pointing out that some other agents had been identified and purified from certain *Paederus* beetles. Pavan [157] seems to be the first who dealt with its biological activity, by working on possible influence of paederin derived from *P. fuscipes* on tumor fragments. He reported that 'paederin' reduced or completely prevented tumor growth and that it acted as an antimitotic, provoking chromosomal alterations and blocking the formation of the spindle at metaphase. Hisada and Emura [158] carried out a cytological study on the antitumor action of paederin, use being made of crude methanol extract from *P. fuscipes*. The extract exerted a considerable influence upon both interphase and mitotic cells, inhibiting mitosis.

Much research has been focused on antimicrobial peptides derived from insect immune defense reactions due to their potential in the development of new antibiotics. Using the gel filtration and reverse-phase high-performance liquid chromatography, Memarpoor-Yazdi et al. [159] identified and purified a new antimicrobial peptide from *Paederus* spp., named sarcotoxin PD which had broad-spectrum inhibitory effects on the examined microbes (bacteria and fungi). The authors suggested the role of sarcotoxin Pd in the treatment of systemic microbial illnesses.

2. Beetles' vesicating products

2.1. Cantharidin

Generally, autogenous producers of cantharidin ($C_{10}H_{12}O_4$) (haemolymph exudation through leg joints or/and antennal pores) occur exclusively within the coleopteran families of Meloidae [13, 48, 161, 162] and Oedemeridae [26, 78, 79, 80, 163, 164] in which it occurs in the haemolymph and other tissues [38, 39, 40, 165]. Cantharidin was discovered in haemolymph and gonads of the blister beetle, commonly known as Spanish fly *Lytta vesicatoria* (Linnaeus), in quantity larger than any other member of blister beetles [166]. It is highly toxic to a wide variety of animals, including birds, amphibians and mammals [13].

It was suggested that cantharidin might be used by the beetle female when selecting a mate at close range [12]. As pointed out in the iron cross blister beetle *Tegrodera aloga* Skinner, individual males donated a large cantharidin-containing spermatophore to their mates, and body size was correlated with the size of the cantharidin-producing accessory glands [167]. Cantharidin production varies widely depending not only on the species but also on the sex, time of year and food source [40] as well as the mating status [48]. The transferring mechanism of some quantity of cantharidin from adult male into the genitalia of female, as a nuptial gift for mating, was clarified by Nikbakhzadeh et al. [168].

Meloid beetles have been attacked by some insects and other natural enemies. It was reported that some blister beetles such as *Meloe* spp. are attacked by *Pedilus* (Coleoptera: Pedilidae) [169] and/or by some species of Miridae (Heteroptera) [170]. When attacked or disturbed, adults of blister beetles release haemolymph droplets in so called "reflex bleeding". The highly toxic material, "cantharidin" in the haemolymph, is a well defensive reaction against the aggressive creatures. As reviewed by Ghoneim [47], cantharidin is considered responsible for the repellent properties of meloid haemolymph against a wide variety of predators.

Although the cantharidin biosynthesis in the blister beetles had been studied by some researchers [48, 171, 172], it was a debatable issue along several decades ago and the mechanism *in vivo* was not completely understood. However, it is quite sure that it implicates reactions with as first compound a terpenic alcohol. A proof can be given with mass spectroscopy using the ^{14}C , 3H , ^{18}O isotopes. The farnesol is a sesquiterpenic alcohol and is an intermediate in the biosynthesis of isoprenoids. Cantharidin was chemically isolated from *M. proscarabaeus* [173].

Cantharidin content was higher in males than in females of the same species. Moreover, cantharidin content in the female collected after the copulation peak was higher than that before copulation peak [174]. Under laboratory conditions, Wang et al. [175] analyzed the changes in the cantharidin level at different life stages of the blister beetle *Mylabris cichorii* Linnaeus. Their results showed that larvae accumulated cantharidin as they grow and develop and adults exhibited a pronounced sexual dimorphism in cantharidin biosynthesis, but total content of cantharidin produced by sex-mixed rearing group was much higher than that by sex-segregated rearing group.

2.2. Paederin

As aforementioned, paederin is a vesicant toxic agent found in the haemolymph of the *Paederus* genus. It differs from cantharidin, in its chemical and biologic properties [26]. The paederin manufacture is largely confined to adult female beetles. Larvae and males only store paederin acquired maternally (i.e., through eggs) or by ingestion [27]. On the other hand, some authors reported only small quantities of the toxin (0.1 to 1.5 μg) in the adult males, while females present up to ten times more paederin (0.2 to 20 μg). Females, themselves, can also be divided into two groups according to the paederin proportion [123]. Positive (+) females predominate in every generation, whereas the negative ones (-) are the minority. So, there is a polymorphism in hemolymph toxin of *Paederus* beetles. Paederin polymorphism is another factor that must be considered in the provocation of linear dermatitis, since positive females constitute most of the *Paederus* population [100]. In the *Paederus* beetles, paederin serves as a chemical defense against predators such as spiders [118] which has not been correctly demonstrated [100].

It has been demonstrated that the paederin is produced not by the beetles themselves but relies on the activities of some species of endosymbiont bacteria within the beetle. According to 16S ribosomal DNA data, only these paederin-

synthesizing females harbor a bacterium with the closest relationship to *Pseudomonas aeruginosa* (Schröter) (Pseudomonadales: Pseudomonadaceae) [176]. Not all species of the genus *Paederus* contain endosymbiotic bacteria and uninfected beetles do not produce paederin [177]. Paederin circulates in the haemolymph of a beetle, so it can be carried over to the offspring (egg, larva, pupa, and beetle). For some details, adult females that are infected with endosymbiotic bacteria produce eggs containing the bacteria in the outer shell walls. Subsequent generations of beetles acquire these endosymbiont bacteria by ingesting some of the 'infected' egg shells or by cannibalizing larvae containing the bacteria [27, 34]. If eggs of such beetles are fed to nonproducing females, paederin is again synthesized in their offspring [178].

Piel et al. [179] had sequenced an extended region of the symbiont genome to gain further insight into the biology of this as-yet-unculturable bacterium and the evolution of paederin symbiosis. Their data indicated that the endosymbiotic bacterium *P. aeruginosa* acquired several foreign genetic elements by horizontal gene transfer. Besides one functional tellurite resistance operon, the region contains a genomic island spanning 71.6 kb that harbors the putative paederin biosynthetic genes. Costa-Sá et al. [180] focused on a search of *Pseudomonas* species and functions of amino acid sequences in the paederin family. They had a better understanding of amino acids paederin cluster and its related functions with the likely *Pseudomonas* species providing that new insights emerge of these genes and proteins to delineate molecular targets for future development of pharmacologically active products.

Chemically, paederin ($C_{25}H_{45}O_9N$) is a toxic amide with two tetrahydropyran rings. It depends on the beetle species and may be one or a combination of several similar molecules including paederone and pseudopaederin [106, 181]. Paederin and several analogues had been prepared through an efficient route that proceeds in ten steps (longest linear sequence) from isobutyraldehyde [182]. Paederin has been synthesized and, in pure form, is a crystalline amide soluble in alcohol and water [100].

Concerning the toxicity, paederin is a potent vesicant toxic, more potent than cobra venom. The toxic effects of paederin were unknown to Western medicine until this century, but they were known to Chinese medicine 1,200 years earlier [106, 114]. It is more toxic externally and internally to rabbits, rats, mice, and guinea pigs than to hedgehogs, chickens, frogs, and toads. It is not known to be toxic to insects, is not attractive to insects, and has little effect against bacteria [183]. It is worthy pointing out that paederin is of natural products with potent antitumor and antiviral activities [184].

3. Prevalence of *Paederus dermatitis* in the world

Although blister beetles are recorded causing human dermatitis, due to their cantharidin [185], meloid or oedemerid dermatitis is not as widespread as PD and remains a minor rural health problem [11]. The dermal effects of paederin from staphylinid beetles have received several names in different countries, but the widely applicable name is linear dermatitis or dermatitis linearis and some of the preferred names are PD and blister beetle dermatitis [108,186]. The irritant contact dermatitis can be caused by the settlement of or accidentally crushed *Paederus* beetles against the skin [187], releasing the paederin [11]. Wet and sticky skin may facilitate local penetration of insect toxins and therefore enhance deleterious effects in use [188,189]. However, *Paederus* species may cause linear lesions [190, 191] and the so-called "kissing lesions", due to the contact between two peculiar skin surfaces, as for example the creases of axillae and elbows [122, 187]. Epidemics caused by *P. sabaeus* is called "bruleuses" in Guinea [191], "skirt and blouse" in Southern Nigeria [190, 192], in Uganda [193], "rove beetle" in South West Africa or the "escaper" in several African countries [142,194], as well as "panchos" in Mexico [195].

PD has been spread in different parts of the world and outbreaks were reported in various countries including Australia [4,196, 197], Japan [198, 199], China [155, 200, 201], Malaysia [3, 125, 126, 202, 203], Korea [204, 205, 207], India [11, 207-211], Sri Lanka [5, 212, 213], Pakistan [214], Iran [34, 103, 128, 215, 216], Iraq [217], sub-Saharan Africa [218], Central Africa [219], Nigeria [190, 220, 221], Tanzania [222], Guinea [191, 223, 224], Uganda [193], Gambia [225], Kenya [217, 226], Sierra Leone [106, 227], Congo [228, 229], Malawi and Namibia [194], South Sudan [230], Egypt [188, 231], Italy [122, 123], France [232, 233, 234], Turkey [202, 235, 236, 237], United States of America [238], Brazil [28], Venezuela [239], Venezuela [240, 241], Mexico [195], Argentina [242], Peru [243], Ecuador [244] and some other countries of Latin America [245].

Skin disorders and dermatitis had been documented among persons in military units or troops in different parts in the world [246, 247, 248, 249]. Christmas et al. [36] described seventy-four New Zealand Army personnel with a distinctive bullous dermatosis caused by an endemic beetle *Thelyphassa lineata* (Fabricius) (Oedemeridae) as the first reported cases. PD was reported in a military unit training in the Arizona desert (USA) during heavy rain and flooding [238]. The first report to an outbreak of PD among U.S. personnel deployed to Pakistan was described. Up to 10% of the base population was afflicted with a pustular eruption and an accompanying halo of erythema [214]. *P. sabaeus* was responsible for a widespread occurrence of PD in United Nations (UN) troops posted in Congo [229].

4. Clinical Symptoms of human dermatitis and conjunctivitis

4.1. Dermatitis

Cantharidin signs and symptoms usually begin within 12 to 24 hours, which is sooner than paederin's effects (Alexander, 1984) while some authors pointed out that a slight redness of the affected areas within the first 12-24 hours is usual in PD [106, 193, 250]. Another difference is that victims usually remember coming into contact with Meloid or Oedemerid beetles because of their large size. Many victims are not aware that *Paederus* were ever crushed on their skin because of their small size and the delayed onset of blistering [82]. In addition, Singh et al. [251] noticed that signs of blister beetle dermatitis practically appeared on most of the patients in early morning after sleep [108] and Padhi et al. [252] reported the same observation.

With regard to the diagnostic features, it is difficult to differentiate PD from other inflammatory skin conditions, without prior experience of managing the infection. The uncommon association of acute dermatitis with minimal or no complaints, which would be noteworthy in the case of chemical or thermal burns, facilitates diagnosis which is corroborated by the season and by the case history [253]. History, predilection for exposed parts, linear nature of primary lesion and presence of kissing lesions are important clue for diagnosis [209]. Capture of the insect, epidemiologic features and a high clinical suspicion can aid in making the correct diagnosis of PD [245].

The common symptoms of PD are skin lesions as well as respiratory, eye or skin allergies that are caused by contact with *Paederus* beetle haemolymph and/or toxic secretions, hairs or frass of larvae [106, 254, 255]. Different responses are seen in the skin depending upon the paederin concentration, duration of exposure, and individual characteristics [250]. In mild cases, there is a slight erythema lasting for a couple days. The erythema is followed by a squamous stage when the blisters dry out over a week, and then a stage when they desquamate, leaving hyper- or hypopigmented patches. The lesions are characteristically linear due to smearing the crushed insect across the skin [118]. As recorded by Assaf et al. [231], blisters caused by the rove beetle *P. alfierii* exhibited a linear configuration in 40% of the patients and kissing lesions were observed in 13%. Multiple lesions occurred in 78% of the patients. The onset of blister formation was reported on the 2-4th day followed by pain [256]. The blister formation may be a response to serine proteases and neutrophil elastase released in response to paederin [257, 258]. Following blister formation there is the appearance of linear lesions (that look like whip injuries) and are also known as mirror image or kissing lesions [239]. Severe cases may demonstrate additional symptoms, such as fever, neuralgia, arthralgia, and vomiting [118, 250].

Concerning the affected parts of the human body, PD symptoms appeared over the exposed areas, usually face, neck and upper extremities [5, 11]. In a study, the most common location of rash was the neck and face (31%) and so hand (30%) but in other studies, the face was the most commonly involved site [231] and the trunk was the least common site [216]. As observed by Ahmad et al. [125] in Malaysia, face and neck were found to be the most frequent sites of PD.

4.2. Conjunctivitis

As previously mentioned, skin blistering by *Paederus* beetles most often occurs on exposed skin of body parts such as the neck, face, arms, and legs where a beetle is felt crawling on the skin and subsequently crushed. On the other hand, paederin is sometimes transferred to other sensitive areas of the body such as eyes or genitals by paederin-contaminated hands [259] as reviewed herein.

Conjunctivitis results from transferring the toxin of crushed beetles by finger to the eyes as a secondary infection [5, 103, 190, 223, 260]. These ocular and periocular lesions in the conjunctivae and eyelids are commonly known as "Nairobi eye" in east Africa [82, 261] caused mainly by the rove beetles *P. sabaeus* [262] and *P. fuscipes* ("spider lick") [263].

Some attention had been paid to the human conjunctivitis as caused by *P. crebipunctatus* [264]. In India, the commonest insect causing Ophthalmic-dermatozoonosis is *P. fuscipes*. In the eye-lids, there may be oedema and tenderness. Occasionally there may be conjunctival hyperemia, chemosis and corneal abrasion [265]. Cases of *Paederus basalis* Bernhauer dermatitis were reported among people of rural areas of Saharanpur and Haridwar districts (India). In the case of infection on eye, odema of eyelid develop and acute conjunctivitis occurs. Eyelids get swollen and eye is closed [210]. Verma and Gupta [266] reported some ocular manifestations due to the rove beetle *P. sabaeus*. Morsy et al. [267] proved experimentally that an extract from *P. alfieri* causes conjunctiva in 75% of used mice. Hashish [188] recognized periorbital dermatosis in people inhabiting the surrounding area of Tanta (Egypt) as caused by *P. alfieri*. He concluded that contact of cornea with paederin from this beetle explained the superficial punctate keratitis and might explain sub-conjunctival hemorrhage or the hemorrhage might be attributed to excessive rubbing of the eye, i.e. mechanical. In a number of patients, clinical findings on the eyelid skin were observed by Canan et al. [268] as erythematous patches and plaques with multiple pustules, erosions and bulla remnants on an erythematous base, squamous and crusted patches and plaques, and erythematous patches. However, conjunctivitis from *Paederus* is rare in the world [269].

5. Mode of action of cantharidin and paederin

Cantharidin is absorbed by the lipid layers of epidermal cell membranes. Application of cantharidin onto the epidermis results in the activation or release of neutral serine proteases that cause degeneration of the desmosomal plaque, leading to the detachment of tonofilaments from desmosomes. This process leads to acantholysis and intra-epidermal blistering, and nonspecific lysis of skin [270]. Lesions heal without scarring, as acantholysis is intra-epidermal [271].

Paederin is a toxic alkaloid, vesicant capable of blocking the mitosis at levels as low as 1 ng/ml, by inhibiting protein and DNA synthesis without affecting RNA synthesis [106, 272], so it prevents the cell division leading to cell death [27]. Histopathological examination of the blister area caused by *P. alfieri* or *P. basalis* revealed features of acute irritant dermatitis in the upper epidermis. Mitotic figures and apoptotic changes such as chromatin condensation and DNA fragmentation were identified in the basal and suprabasal layers [210, 231]. Additionally, foci of acantholysis attributed to the release of epidermal protease have been reported [250]. On the other hand, Caroline [100] hypothesized the transference of endosymbiotic *Pseudomonas* from the beetle to the skin and continues to produce paederin which then leads to the symptom of dermatitis.

6. Preventive measures and curative treatments of dermatitis

Paederus beetles can be prevented by increasing public awareness [273]. They do not enter houses if the windows and doors or ventilation openings are covered with mosquito netting. The beetle usually prefers the bright lights at night, so turning off the lights at night while sleeping or decreasing use of artificial lighting and avoiding fluorescent lights may be useful [230, 245]. If a *Paederus* beetle lands on the skin, it should be blown off and the area in contact should be immediately washed with soap and water [274]. Also, simple preventive measures include an insect repellent cream on exposed parts of the body at night [209]. Ghaffari et al. [275] investigated the repellency effect of the neem oil (*Azadirachta indica* Juss), as a personal protection method against *Paederus* beetles. The results showed that protection produced by Neem oil at 2.5, 5 and 10 % dose was 46.2, 69.3 and 84.62 %, respectively. If the beetles are found on the skin, brushing them off, rather than crushing them, avoids producing PD [230]. The Public Health and Municipal Authorities can control the swarming of rove beetles in residential areas by applying insecticide via thermal fogging or target spraying using Pesguard FG 161 (d-tetramethrin 4.4% w/w/ cyphenothrin 13.2% w/w) glue traps, electric insect traps, etc [100].

Cantharidin-caused skin blisters occur on the neck and arms and other uncovered areas of the body as previously mentioned. General handling of adult beetles seldom results in blistering unless the haemolymph contacts the relatively

thin skin [32]. There is no known antidote for the cantharidin-caused dermatosis [276]. The blistered site should be cleaned with cool wet soaks [259]. Beta-adrenergic agonists and antagonists inhibited the cantharidin dermatitis [277]. Treatment with acetone, ether, fatty soap, or alcohol helps to dissolve and dilute the cantharidin [278]. Iserson [230] suggested rapid washing the affected area, applying cold, wet compresses, and possibly treating with antihistamines.

In East Africa, local remedies (e.g. toothpaste and mud) are used to treat PD, though all are ineffective. The use of cold compressed magnesium sulphate is replacing older remedies [260]. On the basis of its pharmacological properties, Wang [272] applied Moist Exposed Burn Ointment (MEBO) to treat the PD, and it had proved better than traditional methods for its remarkable analgesic, rapid wound-healing and potent anti-infective effects. Nikookar et al. [279] carried out a study to compare the therapeutic effects of atorvastatin for treatment of PD on an experimental group (15 patients) with the control group (15 patients) received placebo. In West Africa, Qadir et al. [227] studied 50 cases of PD at Sierra Leone aiming to determine clinical patterns of PD and its response to topical steroids, with and without antibiotics. They concluded that the better response to a combination of topical steroids, such as betametasone hydrocortisone1% ointment or other anti-inflammatory ointments, and oral antibiotics may indicate concurrent bacterial infection. In Eastern China, the relationship between the toxin of *P. fuscipes* and PD was studied by Wang et al. [155] who observed severe PD and inflammation of eyes in the experimental rabbits caused by the beetle soaked in 70% alcohol for six months. The toxic effect of the beetle on the skin could be inhibited by storing the beetle at 4°C for 6 hours before spreading. In Northern Iran, Davoudi et al. [280] investigated the efficacy of fluocinolone cream, triclocarban (TCC) soap and betamethasone lotion in comparison with placebo in the treatment of PD. They concluded that topical corticosteroids displayed a relative and significant improvement in PD. In Northern Iran, also, Nikookar et al. [279] compared the effectiveness of triamcinolone ointment and atorvastatin tablet with placebo in treatment of PD. The results showed both of triamcinolone ointment and oral atorvastatin had similar effect on PD. Because the PD is a self-limited disease use of topical therapy for treatment of the disease is recommended.

After causing PD in the guinea pig, as an experimental animal, the application of potassium permanganate with calamine to heal PD was more effective than fluocinolone treatment. This topical corticosteroid is thus considered less able to avert the cytotoxic action of paederin on the skin than the antipruritic and cleansing agents [281]. Narendhirakannan et al. [282] assessed a new combination of seven medicinal plants against blisters and infection caused by the rove beetle *P. fuscipes*. Further wound healing property of each plant had been determined which shows Neem with high wound healing power followed by turmeric, *Triphala*, African spider plant and wheatgrass.

The use of steroid topical preparations, pain killers and anti-histamines as beneficial agents had been recommended for the prevention of secondary infections. If there is a secondary infection, removal of irritant, initial washing with soap and water, and application of cold wet compresses followed by steroid and antibiotics should be recommended [5, 103, 187, 190, 209, 223]. Recovery is a slow process and the climatic conditions including high humidity may delay the process of recovery [283]. Recovery occurs within 8-10 days as the lesions dry and dark scales appear [106, 193]. After four weeks the black scales start to peel off, exposing new pink skin. After 6-12 months the scars vanished [283].

References

- [1] Powell JA. Coleoptera. In: Resh VH, Cardé RT (eds.). Encyclopedia of Insects, 2nd ed. Academic Press. **2009**, p. 1132.
- [2] Mumcuoglu Y, Rufli T. Dermatological entomology. 15. Coleoptera/beetles. Schweizerische Rundschau für Medizin (Praxis), 69(47), **1983**, 1750-1755.
- [3] Mokhtar N, Singh R, Ghazali W. Paederus dermatitis among medical students in USM, Kelatan. Med. J. Malaysia, 48, **1993**, 403-406.
- [4] Todd RE, Guthridge SL, Montgomery BL. Evacuation of an Aboriginal community in response to an outbreak of blistering dermatitis induced by a beetle (*Paederus australis*). Med. J. Australia, 164, **1996**, 238-240.
- [5] Kamaladasa SD, Perera WDH, Weeratunge L. An outbreak of paederus dermatitis in a suburban hospital in Sri Lanka. Int. J. Dermatol., 36 (1), **1997**, 34-36.

- [6] Dinev I. The darkling beetle (*Alphitobius diaperinus*)-a health hazard for broiler chicken production. *Trakia J. Sci.*, 1, **2013**, 1-4.
- [7] Ibrahim AA. Infestation with the tiger beetle *Myriochile melancholica* F. (Cicindelidae: Coleoptera) in the student hostel of Benha University, Egypt. *J. Entomol.*, 2(1), **2005**, 59-63.
- [8] Chung RSH, Chew RY, Au-Eong KG. Coleoptera Cincindelidae beetle in the eye. *Singapore Med. J.*, 46(10), **2005**, p.564.
- [9] Rustin MHA, Munro DD. Papular urticaria caused by *Dermestes maculatus* Degeer. *Clin. Exp. Dermatol.* 9, **1984**, 317-321
- [10] Ramachandran S, Hern J, Almeyda J, Main J, Patel KS. Contact dermatitis with cervical lymphadenopathy following exposure to the hide beetle, *Dermestes peruvianus*. *British J. Dermatol.*, 136, **1997**, 943-945.
- [11] Gnanaraj P, Venugopal V, Mozhi MK. An outbreak of *Paederus dermatitis* in a suburban hospital in south India: a report of 123 cases and review of the literature. *J. Am. Acad. Dermatol.*, 57, **2007**, 297-300.
- [12] McCormick JP, Carrel JE. Cantharidin biosynthesis and function in meloid beetles. In: Prestwich GD, Blomquist GJ (eds.). *Pheromone biochemistry*, **1987**, pp: 307-350, Academic Press. Orlando, Florida.
- [13] Dettner K. Inter- and intra-specific transfer of toxic insect compound cantharidin. In: Dettner K, Bauer G, Völkl W (eds.), *Vertical Food Web Interactions*. Springer Verlag. Berlin, **1997**, pp. 115-145.
- [14] Sandroni P. Aphrodisiacs, past and present: a historical review. *J. Clin. Autonomic Res Soc.*, 11(5), **2001**, 303-307.
- [15] Al-Basheer M, Hijazi M, Dama T. Blister beetles dermatosis: a report of 43 cases in a military unit in Eritrea. *J. Royal Med. Serv.*, 9(2), **2002**, 40-43.
- [16] McCluskey A, Keane MA, Walkom CC, Bowyer MC, Sim ATR, Young DJ, Sakoff JA. The first two cantharidin analogues displaying PP1 selectivity. *Bioorg. Med. Chem. Lett.*, 12, **2002**, 391-393.
- [17] Nikbakhtzadeh MR, Tirgari S. Cantharidin component of Iranian blister beetles (Col: Meloidae) and their differences between Iranian and exotic species. *Iranian J. Publ. Health*, 31 (3-4), **2002**, 113-117.
- [18] Bhattacharjee P, Brodell RT. Cantharidin. In Brodell RT, Johnson SM, eds.). *Warts: Diagnosis and Management-an Evidence-Based Approach*. London: Martin Dunitz., **2003**, pp. 151-160.
- [19] An WW, Gong XF, Wang MW, Tashiro SI, Onodera S, Ikejima T. Norcantharidin induces apoptosis in HeLa cells through caspase, MAPK, and mitochondrial pathways. *Acta Pharmacologica Sinica*, 25(11), **2004**, 1502-1508. (In Chinese with English summary).
- [20] Nikbakhtzadeh MR, Ebrahimi B. Detection of cantharidin related compounds in *Mylabris impressa* (Coleoptera: Meloidae). *J. Venomous Animals and Toxins including Tropical Diseases*, 13(3), **2007**, 686-693.
- [21] Zheng LH, Bao YL, Wu Y, Yu CL, Meng XY, Li YX. Cantharidin reverses multidrug resistance of human hepatoma HepG2/ADM cells via down-regulation of P-glycoprotein expression. *Cancer Letters*, 272, **2008**, 102-109.
- [22] Mebs D, Pogoda W, Schneider M, Kauert G. Cantharidin and demethylcantharidin (palasonin) content of blister beetles (Coleoptera: Meloidae) from southern Africa. *Toxicon*, 53(4), **2009**, 466-468.
- [23] Al-Binali AM, Shabana M, Al-Fifi S, Dawood S, Shehri AA, Al-Barki A. Cantharidin poisoning due to blister Beetle ingestion in children: two case reports and a review of clinical presentations. *Sultan Qaboos Univ. Med. J.*, 10(2), **2010**, 258-261.

- [24] Nikbakhtzadeh MR, Vahedi M, Vatandoost H, Mehdinia A. Origin, transfer and distribution of cantharidin-related compounds in the blister beetle *Hycleus scabiosae*. J. Venomous Animals and Toxins including Tropical Diseases, 18 (1), **2012**, 88-96.
- [25] Jiang Y-M, Meng Z-Z, Yue G-X, Chen J-X. Norcantharidin induces HL-60 cells apoptosis *in vitro*. Evidence-Based Complementary and Alternative Medicine, Article ID 154271, **2012**, 4 pp.
- [26] Nicholls DSH, Christmas TI, Greig DE. Oedemerid blister beetle dermatosis: a review. J. Am. Acad. Dermatol., 22(1), **1990**, 815-819.
- [27] Piel J. A polyketide synthase-peptide synthetase gene cluster from an uncultured bacterial symbiont of *Paederus* beetles. Proc. National Acad. Sci., 99 (22), **2002**, 14002-14007.
- [28] Diogenes MJ. Contact dermatitis by pederin: clinical and epidemiological study in Ceara State, Brazil. Rev. Inst. Med. Trop., Sao Paulo, 36(1), **1994**, 59-65. (In Portuguese with English abstract).
- [29] Arnett, R.H. Jr. (2008). False blister beetles, Coleoptera: Oedemeridae. Featured Creatures. EENY-154. http://entomology.ifas.ufl.edu/creatures/urban/medical/false_blisters.htm (19 August 2009).
- [30] Kartal DSP, Atakan D, Eskio glu F. Cantharidin treatment for recalcitrant facial flat warts: a preliminary study. J. Dermatol. Treat., 20(2), **2009**, 114-119.
- [31] Méndez E, Sáenz RE, Johnson CM. Blister dermatitis caused by *Epicauta flagellaria* (Erichson) (Coleoptera: Meloidae) species. Rev. Med. Panama, 14(3), **1989**, 139-144. (In Spanish)
- [32] Selander RB, Fasulo TR. DPI Entomology Circular 268. Publication Number: EENY-166, Copyright 2000-2010 University of Florida, **2010**.
- [33] Mizota K. Additional records on dermatitis caused by three oedemerid species (Coleoptera: Oedemeridae). Med. Entomol. Zool., 52(1), **2001**, 63-66.
- [34] Nikbakhtzadeh MR, Tirgari S. Medically important beetles (Insecta: Coleoptera) of Iran. J. Venomous Animals and Toxins including Tropical Diseases, 14, **2008**, 597-618.
- [35] Abtahi SM, Nikbakhtzadeh MR, Vatandoost H, Mehdinia A, Rahimi-Foroshani A, Shayeghi M. Quantitative characterization of cantharidin in the false blister beetle, *Oedemera podagrariae ventralis*, of the Southern slopes of Mount Elborz, Iranian J. Insect Sci., 12(152), **2012**, 1-5.
- [36] Christmas TI, Nicholls D, Holloway BA, Greig D. Blister beetle dermatosis in New Zealand. New Zealand Med. J., 100(830), **1987**, 515-517.
- [37] Sendur N, Savk E, Karaman G. Paederus dermatitis: a report of 46 cases in Aydin, Turkey. Dermatology, 199, **1999**, 353-355.
- [38] Bologna MA. Coleoptera Meloidae, Fauna d'Italia. XXVIII. (Bologna: Calderini), XIV +, **1991**, 541 pp.
- [39] Bologna MA. A new *Oenas* from Greece and synonymous note on another species from Middle East (Coleoptera, Meloidae). Revue française d'Entomologie, 13(4), **1991**, 175-179.
- [40] Dettner K, Bauer G, Völk W. Vertical Food Web Interactions: Evolutionary Patterns and Driving Forces (Ecological Studies). Springer Verlag. Berlin, **1997**, 412 pp.
- [41] Arnett RH, Thomas MC, Skelly PE, Frank JH. American Beetles, Volume 2. Polyphaga: Scarabaeoidea through Curculionidae. CRC Press, Boca Raton, Florida, **2002**, 861 pp.
- [42] Bologna MA, Pinto JD. The Old World genera of Meloidae (Coleoptera): a key and synopsis. J. Nat. Hist., 36, **2002**, 2013- 2102.

- [43] Bologna MA, Olverio M, Pitzalis M, Mariottini P. Phylogeny and evolutionary history of the blister beetles (Coleoptera, Meloidae). *Molec. Phylogen. Evol.*, 48, **2008**, 679-693.
- [44] Bologna MA, Di Giulio A. Biological and morphological adaptations in the pre-imaginal phases of the beetles family Meloidae. *Atti Accademia Nazionale Italiana di Entomologia, Anno LIX*, **2011**, 141-15.
- [45] Ghoneim KS. Embryonic and postembryonic development of blister beetles (Coleoptera: Meloidae) in the world: A synopsis. *Int. J. Biol. Biol. Sci.*, 2(1), **2013**, 06-18.
- [46] Ghoneim K. Agronomic and biodiversity impacts of the blister beetles (Coleoptera: Meloidae) in the world: A Review. *International J. Agric. Sci. Res.*, 2(2), **2013**, 21-36.
- [47] Ghoneim KS. Behavioral characterization of blister beetles (Coleoptera: Meloidae) in the world: A bibliographic review. *Int. J. Social Behav. Sci.*, 1(2), **2013**, 33-48.
- [48] Carrel JE, McCairel MH, Slagle AJ, Doom JP, Brill J, McCormick JP. Cantharidin production in a blister beetle. *Experientia.*, 49(2), **1993**, 171-174.
- [49] Vazquez-Albalate X. European fauna of Oedemeridae (Coleoptera). *Agraria Editio*, **2002**, pp.178.
- [50] Nikitskyn B. Oedemeridae. In: Ler, P.A. (ed.): A key to the insects of Far Eastern USSR in six volumes. Vol. 3. Coleoptera or Beetles. Part 3. Dalnauka, Vladivostok, **1996**, 09-26. (in Russian).
- [51] Švihla V. A New Genus of the Tribe Asclerini (Coleoptera: Oedemeridae) from Southeastern China. *Ann. Zool.*, 59(4), **2009**, 545-548.
- [52] Mizota K. A New Genus and Species of Asclerini (Coleoptera: Oedemeridae) from the Yaeyama Islands (Japan) and Taiwan. *The Coleopterists' Bull.*, 53, **1999**, 287-292.
- [53] Yoo S, Park HC, Kim JI. Taxonomic review of the subfamily Nacerdinae (Coleoptera: Oedemeridae) in Korea. *Entomol. Res.*, 38(4), **2008**, 287-298.
- [54] Vazquez-Albalate X. Revision of the Southern African Oedemeridae (Coleoptera, Tenebrionoidea). *Mitteilungen aus dem Museum für Naturkunde in Berlin. Zool. Muse. Institut für Spezielle Zool. (Berlin)*, 72(1), **1996**, 83-147.
- [55] Vazquez-Albalate X. Notes on African Oedemeridae, with description of two new species of the genera *Anacerdochroa* and *Probosca* from Southern Africa (Coleoptera, Tenebrionoidea). *Boln. Asoc. esp. Ent.*, 28(3-4), **2004**, 65-79.
- [56] Vazquez-Albalate X. Systematics and phylogeny of the Asclerosibutia-group of genera (Coleoptera, Oedemeridae), with description of a new genus and seven new species from tropical Africa. *Deutsche Entomol. Zeitsch.*, 51, **2004**, 13-52.
- [57] Švihla V. A review of the family Oedemeridae (Coleoptera) of the Socotra Archipelago. *Acta Entomologica Musei Nationalis Pragae*, 52 (suppl. 2), **2012**, 337-346.
- [58] Lencina JL, Gallego D, Andujar C. Nuevos datos de Oedemeridae Latreille, 1810 de la Península Ibérica (Coleoptera). *Heteropterus Rev. Entomol.*, 8(1), **2008**, 95-107.
- [59] Švihla V. Revision of *Chitona* species (Coleoptera: Oedemeridae) from eastern Mediterranean. *Acta Entomol. Musei Nat. Pragae*, 46, **2006**, 107-121.
- [60] Brustel H, Kakiopoulos G. Contribution à la connaissance de la faune de Gréce (Coleoptera: Oedemeridae). *Biocosme Mésogén* (Nice), 26, **2009**, 65-75.

- [61] Barševskis A. Materials on fauna of Oedemeridae (Hexapoda: Coleoptera) in Latvia. 1. *Oedemera subrobusta* (Nakane, 1954) & *Oedemera lurida* (Marsham, 1802). Baltic J. Coleopterol., 9(2), **2009**, 139-150.
- [62] Švihla V. New species of the genus Sparedrus (Coleoptera: Oedemeridae) from Bulgaria. Acta Entomol. Musei Nat. Pragae, 46, **2006**, 123-125.
- [63] Sivilov O. Records of Oedemeridae (Insecta: Coleoptera) species from Strandzha Mountain (Bulgaria and Turkey). ZooNotes, 32, **2012**, 1-5.
- [64] Hojer J. *Oedemera subrobusta* Nakane (*laticollis* auct) an overlooked beetle species in Northern Europe. Entomol. Tidskrift, 129, **2008**, 91-94. (In Swedish with English abstract).
- [65] Švihla V. Revision of the generic classification of the Old World Oedemeridae (Coleoptera). Sborník Národního Muzea v Praze, 41B (1985), **1986**, 141-238.
- [66] Švihla V. Contribution to the knowledge of the palaearctic and oriental Oedemeridae (Coleoptera). Folia Heyrovskyana, 5, **1997**, 83-96.
- [67] Švihla V. Contribution to the knowledge of the Old World Oedemeridae (Coleoptera). Acta Soc. Zool. Bohemicae, 68, **2004**, 61-78.
- [68] Švihla V. Supplementary notes on the family Oedemeridae (Coleoptera) as published in the Catalogue of Palaearctic Coleoptera with the new data of distribution. Klapalekiana, 47, **2011**, 83-88.
- [69] Campbell JM. Family Oedemeridae: false blister beetles. In: Bousquet Y (ed). Checklist of Beetles of Canada and Alaska Publication 1861/E, Agriculture Canada, Research Branch, Ottawa, Ontario, **1991**, 266pp.
- [70] Kriska NL. Oedemeridae Latreille 1810. In: Arnett RH Jr, Thomas MC, Skelley PE, Frank JH (eds.). "American Beetles". Volume 2: Polyphaga: Scarabaeoidea through Curculionoidea. Boca Raton, Florida: CRC Press, **2002**, 514-519.
- [71] Majka CG, Langor D. The Oedemeridae (Coleoptera) of Atlantic Canada. J. Acad. Entomol. Soc., 7, **2011**, 1-6.
- [72] Webster RP, Sweeney JD, DeMerchant I. New Coleoptera records from New Brunswick, Canada: Stenotrachelidae, Oedemeridae, Meloidae, Myceteridae, Boridae, Pythidae, Pyrochroidae, Anthicidae and Aderidae. Zookeys, (179), **2012**, 279-307.
- [73] Blackwelder RE. (Oedemeridae) Checklist of the coleopterous insects of México, Central America, The West Indies, and South America. U.S. National Muse. Bull., 185 (3), **1945**, 489-492.
- [74] Lawrence JF, Newton AF Jr. Families and subfamilies of Coleoptera (with selected genera, notes, references and data on family-group names). In: Pakaluk J, Slipinski SA (eds.). Biology, Phylogeny, and Classification of Coleoptera: Papers Celebrating the 80th Birthday of Roy A. Crowson. Museum i Instytut Zoologii PAN, Warszawa. **1995**, 779-1006.
- [75] Patrice B, Bousquet Y, Davies AE, Alonso-Zarazaga MA, Lawrence JF, Lyal CHC, Newton AF, Reid CAM, Schmitt M, Ślipiński SA, Smith ABT. Family-group names in Coleoptera (Insecta). ZooKeys, 88, **2011**, 1-972 (April, 2011).
- [76] Paula LH. A Systematic revision of the New Zealand Oedemeridae (Coleoptera, Insecta). J. Royal Soc. New Zealand, 5(3), **1975**, 227-274.
- [77] Arnett R.H. Jr. Contribution towards a monograph of the Oedemeridae, 14. A key to and notes on the New World genera. Coleopterists' Bull., 15, **1961**, 49-64.

- [78] Carrel JE, Doom JP, McCormick JP. Identification of cantharidin in false blister beetles (Coleoptera: Oedemeridae) from Florida. *J. Chem. Ecol.*, 12 (3), **1986**, 741-747.
- [79] Frenzel M, Dettner K. Quantification of cantharidin in canthariphilous Ceratopogonidae (Diptera), Anthomyiidae (Diptera) and cantharidin producing Oedemeridae (Coleoptera). *J. Chem. Ecol.*, 20(8), **1994**, 1795-1812.
- [80] Holz C, Streil G, Dettner K, Dütemeyer J, Boland W. Intersexual transfer of a toxic terpenoid during copulation and its paternal allocation to developmental stages: quantification of cantharidin in cantharidin-producing oedemerids (Coleoptera: Oedemeridae) and canthariphilous pyrochroids (Coleoptera: Pyrochroidae). *Zeitschrift für Naturforschung*, 49c, **1994**, 856-864.
- [81] Mehdinia A, Asiabi M, Jabbari A, Abtahi SM. Analysis of cantharidin in false blister beetles (Coleoptera: Oedemeridae) by headspace solid-phase microextraction and gas chromatography-mass spectrometry. *J. Chromatogr, B Analyt. Technol. Biomed. Life Sci.* 1, 879(27), **2011**, 2897-2901.
- [82] Davidson SA, Norton SA, Carder MC, Debboun M. Outbreak of dermatitis linearis caused by *Paederus iliae* and *Paederus iliensis* (Coleoptera: Staphylinidae) at a military base in Iraq. *U.S. Army Med. Depart. J.*, **2009**, 6-15.
- [83] Newton AF. Documenting biodiversity: how well are we doing in Staphyliniformia (Coleoptera). In: the 55th Annual Meet. Entomol. Soc.America. ESA; San Diego, CA. Available to ESA members at: http://esa.confex.com/esa/2007/techprogram/paper_32_168.htm, **2007**.
- [84] Peng Z, Li L-Z, Zhao M-J. Taxonomic study on *Lathrobium Gravenhorst* (Coleoptera, Staphylinidae, Paederinae) from Longwangshan Mountain, East China. *ZooKeys*, 165, **2012**, 21-32.
- [85] Cai C, Huang D. Megolisthaerus, interpreted as staphylinine rove beetle (Coleoptera: Staphylinidae) based on new early cretaceous material from China. *Cretaceous Res.*, 40, **2013**, 207-211.
- [86] Assing V. New species and records of Staphylinidae from Turkey III (Insecta: Coleoptera). *Linzer Biologische Beiträge*, 36 (2), **2004**, 669-733.
- [87] Assing V. New species and records of Staphylinidae from Turkey VI, with six new synonymies (Coleoptera: Staphylinidae). *Koleopterologische Rundschau*, 76, **2006**, 223-276.
- [88] Fagel G. Coleoptera (Staphylinidae): Paederinae. In: Hanstrom B, Brinck P, Rudebeck G (eds). South African animal life, Results of the Lund University Expedition in 1950-1951, 8, **1961**, 259-295. Göteborg: Almqvist & Wiksell.
- [89] Lecoq J-C. Deux nouvelles espèces de Paederinae de Madagascar (Coleoptera, Staphylinidae). *Nouvelle Revue d'Entomologie*, 7 (1), **1990**, 75-80.
- [90] Janák J. Neue Arten neue Funde der Gattung *Oedichirus* aus Madagaskar IV. (Coleoptera: Staphylinidae: Paederinae: Pinophilini). *Klapalekiana*, 39, **2003**, 229-255.
- [91] Oke C. Australian Staphylinidae. *Proc. Royal Soc. Victoria*, 45, **1933**, 101-136.
- [92] Last HR. Records of New Guinea Staphylinidae (Coleoptera) in the Hungarian Natural History Museum. *Ann. Historico-Naturales Muse. Nat. Hungarici*, 72, **1980**, 139-161.
- [93] Herman LH. Revision of *Bledius*. Part IV. Classification of species groups, phylogeny, natural history, and catalogue (Coleoptera, Staphylinidae, Oxytelinae). *Bull. Am. Muse. Nat. Hist.*, 184 (1), **1986**, 1-368.
- [94] Herman LH. Revision of the Asian tribe *Megarthropsini* (Coleoptera: Staphylinidae: Tachyporinae). *Am. Muse. Novitates*, 3430, **2003**, 1-72.

- [95] Herman LH. Generic revision of the Procirrina (Coleoptera: Staphylinidae: Paederinae: Pinophilini). Bull. Am. Muse. Nat. Hist., 347, **2010**, 1-78.
- [96] Herman LH. Revision of the New World species of *Oedichirus* (Coleoptera, Staphylinidae, Paederinae, Pinophilini, Procirrina). Bull. Am. Muse. Nat. Hist., 375, **2013**, 1-137.
- [97] Fernandez LA, Cabrera N, Posse MC, Gabellone C. Catalog of the types of Staphylinidae (Insecta, Coleoptera) deposited at Museo de La Plata (Argentina). Zootaxa, 1611, **2007**, 25-44.
- [98] Brunke A, Klimaszewski J, Anderson RS. Present taxonomic work on Staphylinidae (Coleoptera) in Canada: progress against all odds. ZooKeys, 186, **2012**, 1-5.
- [99] Chatzimanolis S. *Zackfalinus*, a new genus of *Xanthopygina* (Coleoptera: Staphylinidae: Staphylinini) with description of 20 New Species. Ann. Carnegie Muse., 80(4), **2012**, 261-308.
- [100] Caroline A. A review on *Paederus fuscipes* (rove beetle family). Scientia Biologia, 1(1), **2013**, 07-14.
- [101] Dekeirsschieter J, Verheggen FJ, Haubrige E, Brostaux Y. Carrion beetles visiting pig carcasses during early spring in urban, forest and agricultural biotopes of Western Europe. J. Insect Sci., 11, **2011**, 13pp.
- [102] Capinera JL. Dermatitis linearis. Encyclopedia of entomology. Springer, **2008**, 1179 pp.
- [103] Zargari O, Asadi AK, Fathalikhani F, Panahi M. Paederus dermatitis in northern Iran: A report of 156 cases. Int. J. Dermatol., 42, **2003**, 608-612.
- [104] Mullen G, Mullen GR, Durden L. (eds). Medical and Veterinary Entomology. 1st ed., Academic Press, **2002**, 102pp.
- [105] Frank JH. *Paederus sensu lato* (Coleoptera: Staphylinidae): an index and review of the taxa. Insecta Mundi., 2, **1988**, 97-159.
- [106] Frank JH, Kanamitsu K. *Paederus sensu lato* (Coleoptera: Staphylinidae): natural history and medical importance. J. Med. Entomol., 24(2), **1987**, 155-191.
- [107] Abbasipour H, Taghavi A. Study of rove beetles, *Paederus* spp. (Col.: Staphylinidae), the agent of skin dermatitis on north of Iran. In: Lee C-Y, Robinson WH (eds). Proc. 5th Int. Conf. on Urban Pests. Printed by Perniagaan Ph'ng @ P&Y Design Network, Malaysia, **2005**, Abstracts.
- [108] Singh G, Ali S. Paederus dermatitis. Indian J. Dermatol. Venereol. Leprol., 73(1), **2007**, 13-15.
- [109] Nasir S. Biodiversity of staphylinids of Punjab, Pakistan. Ph.D. Thesis. Department of Agri-Entomology, University of Agriculture, Faisalabad, **2011**, 180pp.
- [110] Willers J. Zu einigen paläarktischen Arten der Gattung *Paederus* Fabricius, 1775 (Coleoptera, Staphylinidae, Paederinae). Entomologische Nachrichten und Berichte, 55, **2011**, 255-257. (In German).
- [111] Manley GV. *Paederus fuscipes* (Col.: Staphylinidae): a predator of rice fields in west Malaysia. Entomoph., 22, **1977**, 47-59.
- [112] Mammino JJ. *Paederus* dermatitis: an outbreak on a medical mission boat in the Amazon. J. Clin. Aesthet. Dermatol., 4(11), **2011**, 44-46.
- [113] Min Z. Studies on *Paederus fuscipes* Curtis' artificial diet. Natural Enemies of Insects, 28(1), **2006**, 6-48. (In Chinese with English abstract).
- [114] Nasir S, Akram W. Study of bionomics of skirt and blouse beetle (*Paederus fuscipes* Curtis), 1826 (Coleoptera: Staphylinidae, Paederinae). Pakistan J. Agri. Sci., 49(4), **2012**, 447-450.

- [115] Devi PK, Yadav DN, Anand JHA. Biology of *Paederus fuscipes* Curtis (Coleoptera: Staphylinidae). Pest Manag. Econ. Zool., 10, **2003**, 137-143.
- [116] Laba W, Kilin D. *Paederus fuscipes* Curt. biology and feeding ability of brown planthopper (*Nilaparvata lugens* Stall). Risalah Hasil Penelitian Tanaman Pangan, 4, **1994**, 240-245. (In Indonesian with English abstract).
- [117] Croft A, Smith H, Creamer I. A pseudo-outbreak of skin disease in British troops. J. Royal Soc.Med., 89(10), **1996**, 552-556.
- [118] Kellner RL, Dettner K. Differential efficacy of toxic pederin in deterring potential arthropod predators of *Paederus* (Coleoptera: Staphylinidae) offspring. Oecologia, 107 (3), **1996**, 293-300.
- [119] Kraker J de, Van Huis IA, Van Lenteren JC, Heonge KL, Rabbingea R. Identity and relative importance of egg predators of rice leaffolders (Lepidoptera: Pyralidae). Biol. Control, 19, **2000**, 215-222.
- [120] Ibrahim M. An indication of effect of the wide spread use of some predators in cotton fields. Bull. Soc. Entomol. Egypt, 46, **1962**, 317-323.
- [121] Okiwelu S, Umeozor O, Akpan A. An outbreak of vesicating beetle *Paederus sabeus* Er. (Coleoptera: Staphylinidae) in Rivers State, Nigeria. Ann. Trop. Med. Parasitol., 90(3), **1996**, 345-346.
- [122] Strani GF, Messineo A, Tomidei M. La dermatite da *Paederus fuscipes*. Chron. Dermatol., 13, **1982**, 513-517. (In Italian).
- [123] Veraldi S, Suss L. Dermatitis caused by *Paederus fuscipes* Curt. Int. J. Dermatol., 33(4), **1994**, 277-278.
- [124] Abbasipour H. Bioecology of rove beetle, *Paederus fuscipes* Curtis (Col.: Staphylinidae), causing linear dermatitis in north-west parts of mazandaran province (Iran). In: Lee C-Y, Robinson WH (eds). Proc. 5th Int. Conf. on Urban Pests. Printed by Perniagaan Ph'ng @ P&Y Design Network, Malaysia, **2005**, Abstracts.
- [125] Ahmad A, Dianita R, Abdul-Hamid MW. A study on *Paederus* species (Coleoptera: Staphylinidae) ecology and paederus dermatitis skin disease outbreak in Permatang Pauh, Pulau Pinang. Institut Pengurusan Penyelidikan, Universiti Teknologi Mara, 40450 Shah Alam, Selangor, Malaysia, **2010**, 23pp.
- [126] Bong L-J, Neoh K-B, Jaal Z, Lee C-Y. Life Table of *Paederus fuscipes* (Coleoptera: Staphylinidae). J. Medical Entomol., 49(3), **2012**, 451-460.
- [127] Nasir S, Akram W, Ahmed F. The population dynamics, ecological and seasonal activity of *Paederus fuscipes* Curtis (Staphylinidae; Coleoptera) in the Punjab, Pakistan. Asia-Pacific Chem. Biol. Environ. Engineer. Soc., Procedia, 4, **2012**, 36-41.
- [128] Nikbakhtzadeh MR, Tirgari S. Two medically important beetles of Fars province (Iran): *Paederus* spp. (Col.: Staphylinidae) and its linear dermatitis. Iranian J. Public Health, 28(1-4), **1999**, 73-86.
- [129] Newman DL, Cragg GM, Snader KM. The influence of natural products upon drug discovery. Nat. Prod. Rep., 17, **2000**, 215-234.
- [130] Hengqing JI. Studies for hibernant habitat and hibernant habit of *Paederus fuscipes* Curtis. J. China West Normal Univer. (Nat. Sci.), **2000**, 3-11. (In Chinese with English abstract).
- [131] Judas M, Dornieden K, Strothmann U. Distribution of carabid beetle species at the landscape level. J. Biogeogr., 29, **2002**, 491-508.
- [132] Kehler D, Bondrup-Nielson S, Corkum C. Beetle diversity associated with forest structure including deadwood in softwood and hardwood stands in Nova Scotia. Proc. N.S. Inst.Sci., 42, **2004**, 227-239.

- [133] Aly MZY. Seasonal effect (in Egypt) of moonlight on the vertical distribution of *Paederus alfierii* Koch. (Staphylinidae: Coleoptera). Bull. Entomol. Soc. Egypt, 69, **1990**, 1-9.
- [134] Betz O. Comparative studies on the predatory behavior of *Stenus* spp. (Coleoptera: Staphylinidae): the significance of its specialized labial apparatus. J. Zool., London, 244, **1998**, 527-544.
- [135] Devi PK, Yadav DN, Anand JHA. Predatory behaviour and feeding potential of *Paederus fuscipes*. Indian J. Entomol., 65(3), **2003**, 319-323.
- [136] Berglind SA, Ehnstram B, Ljungberg H. Riparian beetles biodiversity and stream flow regulation- the example of Svartan and Mjallan streams, Central Sweden. Entomologisk-Tidskrift, 118(4), **1997**, 137-154.
- [137] Dash S. Bone marrow metastasis in solid tumours. Indian J. Pathol. Microbiol., 46 (4), **2003**, 613-616.
- [138] Winasa IW, Hindayana D, Santoso S. Release and predation of rove beetle, *Paederus fuscipes* (Coleoptera: Staphylinidae), against the eggs and larvae of *Helicoverpa armigera* (Lepidoptera: Noctuidae) in soybean field. J. Ilmu Pertanian Indonesia, 12(3), **2007**, 147-153.
- [139] Rai MK, Ramamurthy VV, Gupta SL, Dey D. Preliminary investigation on *Paederus fuscipes* (Curtis), a predator in rice agroecosystems. Ann. Plant Protec. Sci., 8(1), **2000**, 2-12.
- [140] Liu K, Zheng F. Functional response of *Paederus fuscipes* Curtis to *Lipaphis erysimi* (Kaltenbach). Journal of Sichuan Teachers College (Natural Science), 25(3), **2004**, 294-297. (In Chinese with English abstract).
- [141] Padmavathi C, Gururaj K, Padmakumari AP, Pasalu IC. Prevalence, prey preference and predatory potential of *Paederus fuscipes* Curtis and *Ophionea* sp. in rice. J. Biol. Control, 22(1), **2008**, 191-193.
- [142] Tawfik M, Kira M, Metwally S. On the abundance of major pests and their associated predators in corn plantations. Bull. Soc. Entomol. Egypt, 58, **1974**, 167-177.
- [143] Tawfik MFS, El-Sherif SI, Abouzied NH. Population fluctuations of the rove-beetle *Paederus alfierii* Koch. (Col., Staphylinidae) in maize, cotton and clover fields in Giza region. Zeitschrift für Angewandte Entomologie, 80(1-4), **1976**, 75-83.
- [144] Shebl MA, Kamel SM, Abu-Hashesh TA, Osman MA. The Most Common Insect Species in Alfalfa Field in Egypt. Acad. J. Entomol., 1 (2), **2008**, 27-31.
- [145] Mesbah H, Saad A, Mourad A, Taman F, Mohamed I. Biological performance of quercetin on the cotton leaf-worm larvae, *Spodoptera littoralis* Boisd. (Lep., Noctuidae) and prevailing natural enemies in the Egyptian cotton fields. Commun. Agric. Appl. Biol. Sci., 72(3), **2007**, 611-622.
- [146] Al-Shannaf HMH. Effect of sequence control sprays on cotton bollworms and side effect on some sucking pests and their associated predators in cotton fields. Egypt. Acad. J. biolog. Sci., 3(1), **2010**, 221- 233.
- [147] El Mandarawy MBR, El Naggar MAZ, Abdel Samae SA. Evaluation of feeding stimulant mixed with chemical and bio-insecticides on certain lepidopterous pests and their natural enemies in cotton and soybean fields. J. Appl. Sci. Environ. Manage., 8(2), **2004**, 77-83.
- [148] Badary H. Ecological aspects of *Sassettia* spp. (Coccidae: Coccoidea: Hemiptera) and their natural enemies in Egypt. Egypt. Acad. J. Biol. Sci., 4 (1), **2010**, 163-174.
- [149] Abd-Rabou S, Ahmed N, Moustafa M. Predators of scale insects (Hemiptera: Coccoidea) and their role in control in Egypt. Egypt. Acad. J. Biolog. Sci., 5(3), **2012**, 203-209.

- [150] Woin N, Ismael S, Bourou S, Bebom T. Potential for biological control of rice yellow mottle virus vectors. African Crop Sci. J., 15(4), **2007**, 211-222.
- [151] Yoshisato I. Effects of several insecticides against *Paederus fuscipes* Curtis (2nd report): LC <-50> values caused by residue contact test. Med. Entomol. Zool., 14 (4), **1963**, 220-222.
- [152] Darwish YA, Farghal AI. Evaluation of certain pesticides activity against the cotton whitefly, *Bemisia tabaci* and associated natural enemies on cotton plants under field conditions in Assiut. Assiut J. Agric. Sci., 21(5), **1990**, 331-339.
- [153] Devi PK, Yadav DN, Jha A. Evaluation of toxicity of some insecticides against *Paederus fuscipes* Curtis (Coleoptera: Staphylinidae). Andhra Agric. J., 48(3-4), **2001**, 263-265.
- [154] Meng Q, Zheng F. Effect of bisultap on predatory function of *Paederus fuscipes* Curtis. Sichuan J. Zool., 24(4), **2005**, 551-553. (In Chinese with English abstract).
- [155] Wang ZX, Shen JL, Xia XF, Wang BT, Wang XH. An experimental observation on the toxin of *Paederus fuscipes* and its causative dermatitis. Acta Parasitol. Med. Entomol. Sinica, 11, **2004**, 107-110 (In Chinese with English summary).
- [156] Fake Z. *Paederus* of causing Dermatitis Linearis and analysis on their beneficial and harmful effects on mankind. J. China West Normal University (Natural Sciences), **1989**, 02. (In Chinese with English summary).
- [157] Pavan M. Ricerche Biologiche e Mediche su Pederina e su Estratti Purificati di *Paederus fuscipes* Curt. (Coleoptera Staphylinidae). Pavia, Impr. Mario Ponzo, Pavia, **1963**, pp. 1-93.
- [158] Hisada Y, Emura M. Cytological Effects of Chemicals on Tumors, XXVIII Notes on the Effect of Extract from *Paederus fuscipes* on a Transplantable Rat Ascites Tumor. J. Fac. Sci. Hokkaido Univ. Ser. VI, Zool. 15, **1965**, 684-692.
- [159] Memarpoor-Yazdi M, Zare-Zardini H, Asoodeh A. A novel antimicrobial peptide derived from the insect *Paederus* dermatitis. Inter. J. Peptide Res. Therap., 19(2), **2013**, 99-108
- [160] Blodgett SL, Carrel JE, Higgins RA. Cantharidin content of blister beetles (Coleoptera: Meloidae) collected from Kansas alfalfa and implications for inducing cantharidiasis. Environ. Entomol., 20, **1991**, 776-780.
- [161] Hemp C, Hemp A, Dettner K. Attraction of the colour beetle species *Pallenothriocera rufimembris* by cantharidin (Coleoptera: Cleridae). Entomol. Gen., 24(1), **1999**, 115-123.
- [162] Hemp C, Hemp A, Dettner K. Canthariphilous insects in east Africa. J. E. Afr. Nat. Hist., 88(1, 2), **1999**, 1-15.
- [163] Carrel JE, Doom JP, McCormick JP. Cantharidin biosynthesis in a blister beetle: inhibition by 6-fluoromevalonate causes chemical disarmament. Experientia, 42(7), **1986**, 853-854.
- [164] Samlaska CP, Samuelson GA, Faran ME, Shparago NI. Blister beetle dermatosis in Hawaii caused by *Thelyphassa apicata* (Fairmaire). Pediatr. Dermatol., 9(3), **1992**, 246-250.
- [165] Young DK. Cantharidin and insects: an historical review. The Great Lakes Entomologist, 17, **1984**, 187-194.
- [166] Bohac V, Winkler JR. A book of beetles. Spring Books, London. **1964**, 150 pp.
- [167] Alcock J, Hadley NF. Tests of alternative hypotheses for assortative mating by size: a comparison of two meloid beetles (Coleoptera: Meloidae). J. Kansas Entomol. Soc., 60(1), **1987**, 41-50.
- [168] Nikbakhtzadeh MR, Dettner K, Boland W, Gäde G, Dötterl S. Intraspecific transfer of cantharidin within selected members of family meloidae (Insecta: Coleoptera). J. Insect Physiol., 53(9), **2007**, 890-899.

- [169] LeSage L, Bousquet Y. A new record of attacks by *Pedilus* (Pedilidae) on *Meloe* (Coleoptera: Meloidae). Entomol. News, 94, **1983**, 95-96.
- [170] Pinto JD. The parasitization of blister Beetles by species of Miridae. Pan-Pacific Entomol., 54, **1978**, 57-60.
- [171] McCormick JP, Carrel JE, Doom JP. Origin of oxygen atoms in cantharidin biosynthesis by beetles. J. Am. Chem. Soc., 108, **1986**, 8071-8074.
- [172] Carrel JE, McCormick JP, Doom JP. Origin of oxygen atoms in cantharidin biosynthesized by blister beetles. Trans. Missouri Acad. Sci., 22, **1988**, 142.
- [173] Dixon AFG, Martin-Smith M, Smith SJ. Isolation of Cantharidin from *Meloe proscarabeus*. Canadian Pharma. J., 29, **1963**, 501-503.
- [174] Yuling F, Juanjie T, Wenzhen M, Jupeng L, Xun L. The natural resource and content of cantharidin in the adult of meloids in China. Acta Entomol. Sinica, 44(2), **2001**, 192-196. (In Chinese with English summary).
- [175] Wang HR, Wang ZK, Chen J, Yang YW, Cao YQ, Xia YX, Yin YP. Changes of cantharidin content in *Mylabris cichorii* Linnaeus (Coleoptera: Meloidae) at different developmental stages under laboratory conditions. Acta Entomol. Sinica, 3, **2008**, 264-268. (In Chinese with English summary).
- [176] Kellner RL. Molecular identification of an endosymbiotic bacterium associated with protein biosynthesis in *Paederus sabaeus* (Coleoptera: Staphylinidae). Insect Biochem. Mol. Biol., 32, **2002**, 389-395.
- [177] Kellner RL. What is the basis of pederin polymorphism in *Paederus riparius* rove beetles? The endosymbiotic hypothesis. Entomol. Exp. Applica., 93, **2003**, 41-49.
- [178] Kellner RLL. Horizontal transmission of biosynthetic capabilities for pederin in *Paederus melanurus* (Coleoptera: Staphylinidae). Chemoecology, 11, **2001**, 127-130.
- [179] Piel J, Höfer I, Hui D. Evidence for a symbiosis island involved in horizontal acquisition of pederin biosynthetic capabilities by the bacterial symbiont of *Paederus fuscipes* beetles. J Bacteriol., 186(5), **2004**, 1280-1286.
- [180] Costa-Sá C, Santos A, Pereira ALC, Diniz MC. Pederin family: identification of symbionts and functions. Resumos do 58º Congresso Brasileiro de Genética, 11-14 de setembro de 2012, Brasil, **2012**. (In Portuguese with English summary).
- [181] Rajesh V, Agarwal S. Blistering beetle dermatitis: An outbreak. Med. J. Armed Forces, India, 62(1), **2006**, 42-44.
- [182] Wu F, Green ME, Floreancig PE. Total Synthesis of Pederin and Analogues. Angewandte Chemie Int. Ed., 50(5), **2011**, 1131-1134.
- [183] Piper R. Extraordinary Animals: An Encyclopedia of Curious and Unusual Animals. Greenwood Publishing Group, Amazon.com, **2007**, 298pp.
- [184] Narquian R, Kocienski PJ. The pederin family of antitumor agents: structures, synthesis and biological activity, In: Mulzer R, Bohlmann R (eds.). The role of natural products in drug discovery, vol. 32. Springer-Verlag, Heidelberg, Germany, **2000**, 25-56.
- [185] Inanir I. Erythema multiforme associated with blister beetle dermatitis. Contact Dermatitis, 46(3), **2002**, 175.
- [186] Peschke K, Friedrich P, Gantert C, Metzler M. The use of the tergal gland defensive secretion in male intrasexual aggression of the rove beetle, *Aleochara curtula* (Coleoptera: Staphylinidae), measured by closed-loop-stripping-analyses and tandem bioassay-mass fragmentography. Chemoecology, 7, **1996**, 24-33.

- [187] Narasimhalu CR, Murali A, Kannan R, Srinivasan N. Blister beetle dermatitis. J. Indian Med. Assoc., 108(11), **2010**, 781-782.
- [188] Hashish AA. Periorbital Paederus Dermatosis: a missed diagnosis. AIOS, 1(3), **2010**, 270-272. Available in www.ios.org.
- [189] Sinan A, Çevik IE. Faunistic studies on Paederinae (Coleoptera: Staphylinidae) in Manisa province, Turkey. Munis Entomol. Zool., 3 (2), **2011**, 665-674.
- [190] George AO, Hart PD. Outbreak of *Paederus dermatitis* in southern Nigeria: epidemiology and dermatology. Int. J. Dermatol., 29, **1990**, 500-501
- [191] Pampiglione S, Trentini M. Dermatite eritemato-vesicolare da *Paederus sabaeus* Erichson 1840 (Coleoptera, Staphylinidae) in Repubblica di Guinea. Ann. Ital. Dermatol. Clin., 45, **1991**, 15-20. (In French).
- [192] George AO, Falope ZF. An epidemic of Paederus dermatitis in southern Nigeria. Contact Dermatitis, 20(4), **1989**, 314-315.
- [193] McCrae AWR, Visser SA. *Paederus* (Coleoptera: Staphylinidae) in Uganda. I. Outbreaks, clinical effects, extraction and bioassay of vesicating toxin. Ann. Trop. Med. Parasitol., 69, **1975**, 109-120.
- [194] Deneys JB, Zumpt F. Rove beetle dermatitis in south west Africa. S. Afr. Med. J., 37, **1963**, 1284-1285.
- [195] Navarrete-Heredia JL, Flores VHG. Aspectos etnoentomológicos acerca de *Paederus* sp. (Coleoptera: Staphylinidae) en Mascota, Jalisco, México. Dugesiana, 12(1), **2005**, 9-18. (In Spanish with English abstract).
- [196] Banney LA, Wood DJ, Francis GD. Whiplash rove beetle dermatitis in central Queensland. Aust. J. Dermatol., 41(3), **2000**, 162-167.
- [197] Peters P, Davies D. Paederus dermatitis in mine workers from insects attracted to high power artificial lighting. Australasian J. Dermatol., 53(1), **2012**, 46.
- [198] Armstrong RK, Winfield JL. Staphylinidae dermatitis on Okinawa. J. Med. Entomol., 5, **1968**, 362.
- [199] Armstrong RK, Winfield JL. *Paederus fuscipes* dermatitis: an epidemic on Okinawa. Am. J. Trop. Med. Hyg., 18, **1969**, 147-150.
- [200] Li SR. A study on investigation of an outbreak of Paederus dermatitis in the city of Yantai. Zhonghua Yu Fang Yi Xue Za Zhi., 24(3), **1990**, 149-50. (In Chinese with English summary).
- [201] Sheng D, Sheng XY. An investigation of an epidemic of *Paederus fuscipes* dermatitis in Guizhou. Chin. J. Dermatol., 28, **1995**, 107-108.
- [202] Rahmah E, Norjaiza MJ. An outbreak of Paederus dermatitis in a primary school, Terengganu, Malaysia. Malays. J. Pathol., 30(1), **2008**, 53-56.
- [203] Heo CC, Latif B, Hafiz WM, Zhou HZ. Dermatitis caused by *Paederus fuscipes* Curtis 1840 (Coleoptera: Staphylinidae) in student hostels in Selangor, Malaysia. Southeast Asian J. Trop. Med. Public Health, 44(2), **2013**, 197.
- [204] Kim S, Park JH, Lee JH, Yang JM, Lee ES, Kim JK, Kim WS. A case of Paederus Dermatitis. Ann. Dermatol., 19(2), **2007**, 88-90. (In Korean with English summary).
- [205] Kwon OC, Lee JY, Lee IY, Cho BK. A Case of Paederus Dermatitis due to paederus fuscipes. Korean J. Dermatol., 37(2), **1999**, 281-283. (In Korean with English abstract).

- [206] You DO, Kang JD, Youn NH, Park SD. Bullous contact dermatitis caused by self-applied crushed *Paederus fuscipes* for the treatment of vitiligo. Cutis, 72(5), **2003**, 385-388.
- [207] Sengupta SR, Lahiri K. Blister beetle dermatitis in West Bengal. Indian J. Dermatol. Venereol. Leprol, 63, **1997**, 69-70.
- [208] Verma R, Agarwal S. Blistering beetle dermatitis: an outbreak. Med. J. Armed Forces, India, 62(1), **2006**, 42-44.
- [209] Kumar P, Gharami RC, Mondal A, Ghosh K. Sudden appearance of erythematous papules and vesicles. J. Pakistan Assoc. Dermatologists, 20, **2010**, 57-59.
- [210] Dhiman SC. *Paederus basalis* Bernhauer dermatitis, its cause and preventive measures. J. Exp. Zool. India, 14(2), **2011**, 485-491.
- [211] Malvankar DD, Sacchidanand S, Mallikarjun M, Nataraj HV. Linear lesions in dermatology. Indian J. Dermatol. Venereol. Leprol., 77, **2011**, 722-726.
- [212] Brazzelli V, Martinoli S, Prestinari F. Staphylinid blister beetle dermatitis. Contact dermatitis, 46(3), **2002**, 183-184.
- [213] Senanayake MP, Jayasena A, Varathan V, Kamaladeepan S. Blister beetle dermatitis: A seasonal occurrence. Sri Lanka J. Child. Health, 40, **2011**, 34.
- [214] Dursteler BB, Nyquist RA. Outbreak of rove beetle (Staphylinidae) pustular contact dermatitis in Pakistan among deployed U.S. personnel. Mil. Med., 169, **2004**, 57-60.
- [215] Davoodi SM, Bakhtiyari P, Khoobdel M. Determination of Sings and symptoms of Paederus dermatitis in Behshahr hospital in 2000. J. military Med., 10(1), **2000**, 29-34. (In Persian with English summary).
- [216] Hajheydari Z. An investigation of clinical and demographic features of paederus dermatitis in Sari dermatological clinics form March to October 2002 . J. Mazandaran Univ. Med. Sci., 14, **2004**, 97-104. (In Persian with English summary).
- [217] Al-Dhalimi MA. Paederus dermatitis in Najaf province of Iraq. Saudi Med. J., 29, **2008**, 1490-1493.
- [218] Krippner R, Dzikouk G. Toxische Kontaktdermatitis durch *Paederus sabaeus* (Blasenkäfer) Eine Epidemie in Afrika südlich der Sahara. Deutsches Ärzteblatt, 97, **2000**, 1072-1074. (In German with English summary).
- [219] Penchenier L, Mouchet J, Cros B, Legall P, Cosnefroy JY, Quézédé P, Chandenier J. Invasions de Paederus sabaeus (Coleoptera: Staphylinidae) en Afrique Centrale. I. Aspects entomologiques et épidémiologiques. Bull. Soc. Path. Ex., 87, **1994**, 45-48. (In French).
- [220] George AO. Paederus dermatitis-a mimic. Contact Dermatitis, 29, **1993**, 212-213.
- [221] Oyedeji OA, Adebami OJ, Oyedeji GA. Linear Dermatitis following contact with the rove beetle in Nigerians: a report of 2 cases. Int. J. Trop. Med., 3(1), **2006**, 6.
- [222] Poole TRG. Blister beetle periorbital dermatitis and keratoconjunctivitis in Tanzania. Eye, 12, **1998**, 883-885.
- [223] Couppie P, Beau F, Grosshans E. Paederus dermatitis: apropos of an outbreak in Conakry (Guinea) in November 1989. Ann. Dermatol. Venereol., 119, **1992**, 191-195.
- [224] Vanhecke C, Malvy D, Guevart E, Laloge V, Ezzedine K. Paederus dermatitis: A retrospective study of 74 cases occurring in 2008 in Guinea-Conakry. Ann. Dermatol. Venereol., 137(3), **2010**, 189-193.
- [225] Giglioli MEC. Some observations on blister beetles, family Meloidae, in Gambia, West Africa. Trans. Royal Soc. Trop. Med. Hyg., 59(6), **1965**, 657-663.

- [226] Schayk IMCJ van, Agwanda RO, Githure JI, Beier JC, Knols BGJ. El Niño causes dramatic outbreak of Paederus dermatitis in East Africa. In: Low, P.S. (ed.). Climate change and Africa. **2005**, 240-247, Cambridge University Press.
- [227] Qadir SN, Raza N, Rahman SB. Paederus dermatitis in Sierra Leone. Dermatol. Online J., 12(7), **2006**, 9.
- [228] Chandenier J, Quézédé P, Chandenier B, Penchenier L, Gathisé A, Mouchet J. Invasions of *Paederus sabaeus* (Coleoptera Staphylinidae) in central Africa. 2. Clinical and therapeutic aspects in Brazzaville. Bull Soc Pathol Exot., 87(1), **1994**, 49-51. (In French)
- [229] Vasudevan B, Joshi DC. Irritant dermatitis to staphylinid beetle in Indian troops in Congo. Med. J. Armed Forces, India, 66(2), **2010**, 121-124.
- [230] Iserson KV. Nairobi Fly (*Paederus*) Dermatitis in South Sudan: A case report. Wilderness & Environmental Medicine, 23(3), **2012**, 251-254.
- [231] Assaf M, Nofal E, Nofal A, Assar O, Azmy A. Paederus dermatitis in Egypt: a clinicopathological and ultrastructural study. J. Euro. Acad. Dermatol. Venereol., 24(10), **2010**, 1197-1201.
- [232] Handa F, Pradeep S, Sudarshan G. Beetle dermatitis in Punjab. Indian J. Dermatol. Venerol. Leprol., 51, **1985**, 208-2012.
- [233] Kalla G, Batra A. Blister beetle dermatitis. Indian J. Dermatol. Venerol. Leprol., 62, **1996**, 267-268.
- [234] Drouet G, Glaizal M, Schmitt C, Kervégant M, Tichadou L, de Haro L. Paederus dermatitis: Four cases in Provence, Southern France. Presse Med., 42(3), **2013**, 355-357. (In French).
- [235] Uslular C, Kavukcu H. An epidemicity of *Paederus* species in the Cukurova region. Cutis., 69, **2002**, 277-279.
- [236] Sendur N, Savk E, Karaman G. Paederus dermatitis: a report of 46 cases in Aydin, Turkey. Dermatology, 199, **2003**, 353-55.
- [237] Erdogan B\$, Ergin \$, Aktan \$, Kaçar N. The cases with Paederus Dermatitis from Denizli Region. Turkderm, 40(4), **2006**, 123-125. (In Turkish with English summary).
- [238] Claborn DM, Polo JM, Olson PE, Earhart KC, Sherman SS. Staphylinid (rove) beetle dermatitis outbreak in the American southwest? Mil. Med., 164(3), **1999**, 209-213.
- [239] Kerdel-Vegas F, Goihman-Yahr M. Paederus dermatitis. Arch. Dermatol., 94, **1966**, 175-185.
- [240] Dao L. Dermatitis caused by Coleoptera Paederus columbines in Venezuela. Dermatol. Trop. Ecol. Geogr., 30, **1964**, 169-72. (In Spanish).
- [241] Rivas LG, Parra JJ, Flores OR. Paederus Dermatitis in the State of Cojedes, Venezuela. Derm. Venez., 39, **2001**, 93-98.
- [242] Sujit SR, Koushik L. Blister beetle dermatitis in West Bengal. Indian J. Dermatol. Venereol. Leprol., 63, **1997**, 69-70.
- [243] Alva-Davalos V, Laguna-Torres VA, Huaman A, Olivos R, Chavez M, Garcia C, Mendoza N. Epidemic dermatitis by *Paederus irritans* in Piura, Perú at 1999, related to El Niño phenomenon. Rev. Soc. Bras. Med. Trop., 35(1), **2002**, 23-28.
- [244] Senel E, Sahin C. A warmer world means more beetles and more dermatitis. Indian J. Occup. Environ. Med. Jan-Apr; 15(1), **2011**, 47.

- [245] Cressey BD, Paniz-Mondolfi AE, Rodriguez-Morales AJ, Ayala JM, Silva AA. Dermatitis Linearis: vesicating dermatosis caused by *Paederus* species (Coleoptera: Staphylinidae): case series and review. *Wilderness & Environmental Medicine*, **2013** (In press).
- [246] Oumeish OY, Rushaidat QM. Contact dermatitis to military boots in Jordan. *Contact Dermatitis*, **6**(7), **1981**, 498.
- [247] Goon ATJ, Goh CL. Occupational skin diseases in national servicemen and military personnel in Singapore, 1989-1999. *Contact Dermatitis*, **44**(2), **2001**, 89-90.
- [248] Wolf R, Orion E, Matz H. Contact dermatitis in military personnel. *Clin. Dermatol.*, **20**(4), **2002**, 439-444.
- [249] Dampierre H. "Epidemic" of plantar dermatitis during military maneuvers in the rain forest of Guyana. *Med. Trop.*, **63**(2), **2003**, 179-182. (In French).
- [250] Borroni G, Brazzelli V, Rosso R, Pavan M. *Paederus fuscipes* dermatitis: a histopathological study. *Am. J. Dermatopath.*, **13**, **1991**, 467-474.
- [251] Singh A, Singh AK, Singh D, Varghese A. Blister beetle dermatitis: few observations helping in diagnosis. *Int. J. Prev. Med.*, **4**(2), **2013**, 241.
- [252] Padhi T, Mohanty P, Jena S, Sirka CS, Mishra S. Clinicoepidemiological profile of 590 cases of beetle dermatitis in western Orissa. *Indian J. Dermatol. Venereol. Leprol.*, **73**, **2007**, 333-335.
- [253] Gelmetti C, Grimalt R. Paederus dermatitis: An easy diagnosable but misdiagnosed eruption. *Euro. J. Pediatrics*, **152**(1), **1993**, 6-8.
- [254] Ahmed AR, Moy R, Barr AR, Price Z. Carpet beetle dermatitis. *J. Am. Acad. Dermatol.*, **5**, **1981**, 428-432.
- [255] Alexander JO. Arthropods and Human Skin. New York: Springer-Verlag, London, **1984**, 422pp.
- [256] Southcott RV. Injuries from Coleoptera. *Med. J. Aust.*, **151**(11-12), **1989**, 654-659.
- [257] Liu Z, Shapiro SD, Zhou X, Twining SS, Senior RM, Giudice GJ, Fairley JA, Diaz LA. A critical role for neutrophil elastase in experimental bullous pemphigoid. *J. Clin. Invest.*, **105**, **2000**, 113-123.
- [258] Liu Z, Zhou X, Shapiro SD, Shipley JM, Twining SS, Diaz LA, Senior RM, Werb Z. The serpin alpha1-proteinase inhibitor is a critical substrate for gelatinase B/MMP-9 *in vivo*. *Cell*, **102**, **2000**, 647-655.
- [259] Rahman S. Paederus dermatitis in Sierra Leone. *Dermatol. Online J.*, **12**, **2006**, 9.
- [260] Fox R. *Paederus* (Nairobi Fly) vesicular dermatitis in Tanzania. *Trop. Doct.*, **23**(1), **1993**, 17-19.
- [261] Williams AN. Rove beetle blistering--Nairobi eye. *J. R. Army Med. Corps.*, **139**, **1993**, 17-19.
- [262] McCrae AWR, Visser SA. Recent "Nairobi eye" beetle (*Paederus sabaeus* Er.) outbreaks in Uganda, and studies on the vesicating toxin. *Biochem. J.*, **89**, **1963**, 79.
- [263] Somerset EJ. "Spider lick", an epidemic ophthalmic-dermatozoonosis due to beetles of the genus *Paederus*. *Brit. J. Ophthal.*, **45**, **1961**, 395-407.
- [264] Roberts J, Tonking H. Notes on an East African vesicant beetle. *Paederus crebipunctatus*. *Ann. Trop. Med. Parasit.*, **29**, **1935**, 415.
- [265] Mukherjee S, Ahmed E. A few observations on cases of ophthalmic-dermatozoonosis. *Indian J. Ophthalmol.*, **14**, **1966**, 178-81.

- [266] Verma S, Gupta S. Ocular manifestations due to econda (*Paederus sabaeus*). Med. J. Armed Forces, India, 68(3), 2012, 245-248.
- [267] Morsy T, Arafa M, Younis T, Mahmoud I. Studies on *Paederus alfieri* Koch (Coleoptera: Staphylinidae) with special references to the medical importance. J. Egypt. Soc. Parasitol., 26(2), 1996, 337-335.
- [268] Canan H, Altan-Yaycioglu R, Durdu M. Periocular Paederus dermatitis mimicking preseptal cellulitis. Can. J. Ophthalmol., 48(2), 2013, 121-125.
- [269] Mbonile L. Acute haemorrhagic conjunctivitis epidemics and outbreaks of *Paederus* spp. keratoconjunctivitis ('Nairobi red eyes') and dermatitis. S. Afr. Med. J., 101, 2011, 541-543.
- [270] Bertaux B, Prost C, Heslan M, Dubertret L. Cantharide acantholysis: endogenous protease activation leading to desmosomal plaque dissolution. Br. J. Dermatol., 188, 1988, 157-165.
- [271] Moed L, Shwayder TA, Chang MW. Cantharidin Revisited: a blistering defense of an ancient medicine. Arch. Dermatol., 137, 2001, 1357-1360.
- [272] Wang X. Clinical experience with MEBO in treating Paederus Dermatitis. Chinese J. Burns Wounds and Surface Ulcers, 2(4), 2000, 60-61.
- [273] Huang C, Liu Y, Yang J, Tian J, Yang L, Zhang J, Li Y, Li J, Wang C, Tu Y, Tao J. An outbreak of 268 cases of Paederus dermatitis in a toy-building factory in central China. Int. Soc. Dermatol., 48, 2009, 128-131.
- [274] Schofield S. Bugs don't have to bite to do damage: the tale of the *Paederus* beetle. 2004. www.forces.gc.ca [Canadian Forces Health Services website]. <http://www.forces.gc.ca/health-sante/wn-qn/adv-avi/Paederus-eng.asp#tphp>.
- [275] Ghaffari D, Tirgari S, Moosavi Ivanaki A. Repellency effect of neem oil (*Azadirachta indica* Juss) against *Paederus* beetles, the agent of linear dermatitis, (Col: Staphylinidae). Modares J. Med. Sci., Pathol., 9, 2007, 45-51.
- [276] Mallari RQ, Saif M, Elbualy MS, Sapru A. Ingestion of a blister beetle (Meloidae family). Pediatrics, 98(3 Pt 1), 1996, 458-459.
- [277] Swingle KF, Reiter MJ, Schwartzmiller DH. Comparison of croton oil and cantharidin induced inflammations of the mouse ear and their modification by topically applied drugs. Arch. Int. Pharmacodyn. Ther., 254(1), 1981, 168-176.
- [278] Burnett JW, Calton GJ, Morgan RJ. Blister beetles: "Spanish fly". Cutis., 40, 1987, 22.
- [279] Nikookar SH, Hajheydari Z, Moosa-Kazemi SH, Mahmoudi M, Shahmohammadi S. Comparison of topical triamcinolone and oral atorvastatin in treatment of paederus dermatitis Northern Iran. Pakistan J. Biol. Sci., 15(2), 2012, 103-107.
- [280] Davoudi SM, Rostami P, Emadi SN, Sadr B, Khabiri E. Comparison of the efficacy of Fluocinolone cream, Triclocarban soap and Betamethasone lotion with placebo in the treatment of Paederus Dermatitis. Iranian J. Dermatol., 9 (37), 2006, 217-220.
- [281] Fakoorziba MR, Eghbal F, Azizi K, Moemenbellah-Fard MD. Treatment outcome of Paederus dermatitis due to rove beetles (Coleoptera: Staphylinidae) on guinea pigs. Trop. Biomed., 28(2), 2011, 418-424.
- [282] Narendhirakannan RT, Nirmala JG, Caroline A, Lincy S, Saj M, Durai D. Evaluation of antibacterial, antioxidant and wound healing properties of seven traditional medicinal plants from India in experimental animals. Asian Pacific J. Trop. Biomed., 2(3), 2012, 245-253.

- [283] Khan TA, Ali I, Singh VV, Sharma S. Utilization of fly ash as low-cost adsorbent for the removal of methylene blue, malachite green and rhodamine B dyes from textile wastewater. *J. Environ. Protect. Sci.*, 3, **2009**, 11-22.